


<http://www.cas.org/support/stngen/stdoc/properties.html>

=>

Uploading C:\Program Files\Stnexp\Queries\10588532.str



chain nodes :

7 8 9 10 11

ring nodes :

1 2 3 4 5 6 12 13 14 15 16 17

chain bonds :

6-7 7-8 8-9 9-10 10-11 10-12

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6 12-13 12-17 13-14 14-15 15-16 16-17

exact/norm bonds :

8-9 9-10 10-11

exact bonds :

6-7 7-8 10-12

normalized bonds :

1-2 1-6 2-3 3-4 4-5 5-6 12-13 12-17 13-14 14-15 15-16 16-17

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:CLASS 8:CLASS

9:CLASS 10:CLASS 11:CLASS 12:Atom 13:Atom 14:Atom 15:Atom


16:Atom 17:Atom

L1 STRUCTURE UPLOADED

=> d l1

L1 HAS NO ANSWERS

L1 STR



Structure attributes must be viewed using STN Express query preparation.

=> s 11 sss full

FULL SEARCH INITIATED 15:07:52 FILE 'REGISTRY'  
FULL SCREEN SEARCH COMPLETED - 18980 TO ITERATE

100.0% PROCESSED 18980 ITERATIONS

3537

ANSWERS

SEARCH TIME: 00.00.01

L2 3537 SEA SSS FUL L1

=> scan


ENTER TERM OR (END):scan 12

ENTER FIELD CODE (BI):bi

E1	2	SCAMPHORSULFONATE/BI
E2	47	SCAN/BI
E3	0 -->	SCAN L2/BI
E4	1	SCANA/BI
E5	1	SCANADI/BI
E6	1	SCANADIUM/BI
E7	1	SCANAL/BI
E8	1	SCANALKA/BI
E9	2	SCANALLOY/BI
E10	1	SCANCEM/BI
E11	1478	SCAND/BI
E12	8	SCAND1/BI

=> d scan


L2 3537 ANSWERS REGISTRY COPYRIGHT 2009 ACS on STN  
IN Benzamide, 2-(aminosulfonyl)-N-[2-(2-pyridinyl)ethyl]-  
MF C14 H15 N3 O3 S



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L2 3537 ANSWERS REGISTRY COPYRIGHT 2009 ACS on STN  
IN INDEX NAME NOT YET ASSIGNED  
MF C26 H28 F N3 O4



HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):0

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s l2 and (anti!fung? or fungicid? or pesticid?)

444 L2

3 ANTI!FUNG?

118517 FUNGICID?

98639 PESTICID?

L3 54 L2 AND (ANTI!FUNG? OR FUNGICID? OR PESTICID?)

=> s l3 and (py<2003 or ay<2003 or pry<2003)

22983274 PY<2003

4503738 AY<2003

3972615 PRY<2003

L4 9 L3 AND (PY<2003 OR AY<2003 OR PRY<2003)

=> d l4 ibib abs 1-9

L4 ANSWER 1 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2004:162541 CAPLUS [Full-text](#)

DOCUMENT NUMBER: 140:176744

TITLE: Preparation of 2-pyridylethylbenzamide derivative

fungicides

INVENTOR(S): Mansfield, Darren James; Cooke, Tracey; Thomas, Peter

Stanley; Coqueron, Pierre-Yves; Vors, Jean-

Pierre;

Briggs, Geoffrey Gower; Lachaise, Helene;

Rieck,

Heiko; Desbordes, Philippe; Grosjean-

Cournoyer,

Marie-Claire

PATENT ASSIGNEE(S): Bayer Cropscience S. A., Fr.


SOURCE: PCT Int. Appl., 35 pp.

DOCUMENT TYPE:  
 LANGUAGE:  
 FAMILY ACC. NUM. COUNT:  
 PATENT INFORMATION:

CODEN: PIXXD2  
 Patent  
 English  
 2

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	-----	-----	-----	-----
WO 2004016088	A2	20040226	WO 2003-EP9516	
20030808 <--				
WO 2004016088	A3	20040325		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
EP 1389614	A1	20040218	EP 2002-356159	
20020812 <--				
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
CA 2492173	A1	20040226	CA 2003-2492173	
20030808 <--				
AU 2003266316	A1	20040303	AU 2003-266316	
20030808 <--				
AU 2003266316	B2	20071025		
EP 1531673	A2	20050525	EP 2003-787805	
20030808 <--				
EP 1531673	B1	20060104		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
BR 2003013340	A	20050712	BR 2003-13340	
20030808 <--				
CN 1674784	A	20050928	CN 2003-819471	
20030808 <--				
CN 1319946	C	20070606		
JP 2005535714	T	20051124	JP 2004-528509	
20030808 <--				
NZ 537608	A	20060428	NZ 2003-537608	
20030808 <--				
RU 2316548	C2	20080210	RU 2005-106877	

20030808 <--  
 IN 2005DN00120 A 20090130 IN 2005-DN120  
 20050113 <--  
 MX 2005001580 A 20050425 MX 2005-1580  
 20050209 <--  
 US 20050234110 A1 20051020 US 2005-524345  
 20050211 <--  
 KR 853967 B1 20080825 KR 2005-702419  
 20050212 <--  
 HK 1080329 A1 20071109 HK 2006-100382  
 20060110 <--  
 PRIORITY APPLN. INFO.: EP 2002-356159 A  
 20020812 <--  
 FR 2003-5233 A  
 20030429  
 WO 2003-EP9516 W  
 20030808  
 OTHER SOURCE(S): MARPAT 140:176744  
 GI



AB The 2-pyridylethylbenzamide derivs. I, in which p is 1, 2, 3 or 4; q is 1, 2, 3, 4 or 5; X is chosen, halo, alkyl or haloalkyl, at least one of the substituents being a haloalkyl; Y is halo, alkyl, alkenyl, alkynyl, haloalkyl, alkoxy, amino, phenoxy, alkylthio, dialkylamino, acyl, cyano, ester, hydroxy, aminoalkyl, benzyl, haloalkoxy, halosulfonyl, halothioalkyl, alkoxyalkenyl, alkylsulfonamide, nitro, alkylsulfonyl, phenylsulfonyl or benzylsulfonyl; as well as I N-oxides are prepared as fungicides. N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-ethyl}- 2,6-dichlorobenzamide is an exception. Method for treating phytopathogenic diseases.

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS

RE FORMAT RECORD. ALL CITATIONS AVAILABLE IN THE

L4 ANSWER 2 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 2004:136478 CAPLUS Full-text  
 DOCUMENT NUMBER: 140:181332  
 TITLE: Preparation of N-{2-(2-pyridyl)ethyl}benzamides as

fungicides  
 INVENTOR(S): Mansfield, Darren James; Cooke, Tracey;  
 Thomas, Peter  
 Stanley; Vors, Jean-Pierre; Coqueron, Pierre-

Yves;

PATENT ASSIGNEE(S):

SOURCE:

DOCUMENT TYPE:

LANGUAGE:

FAMILY ACC. NUM. COUNT:

PATENT INFORMATION:

Briggs, Geoffrey Gower; Lachaise, Helene

Bayer Cropscience S.A., Fr.

Eur. Pat. Appl., 17 pp.

CODEN: EPXXDW



Patent

French

2

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	----
EP 1389614	A1	20040218	EP 2002-356159	
20020812 <--				
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
CA 2492173	A1	20040226	CA 2003-2492173	
20030808 <--				
WO 2004016088	A2	20040226	WO 2003-EP9516	
20030808 <--				
WO 2004016088	A3	20040325		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,				
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,				
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,				
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM,				
PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN,				
TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RN: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,				
KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES,				
FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR,				
BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2003266316	A1	20040303	AU 2003-266316	
20030808 <--				
AU 2003266316	B2	20071025		
EP 1531673	A2	20050525	EP 2003-787805	
20030808 <--				
EP 1531673	B1	20060104		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
BR 2003013340	A	20050712	BR 2003-13340	
20030808 <--				
CN 1674784	A	20050928	CN 2003-819471	
20030808 <--				
CN 1319946	C	20070606		
JP 2005535714	T	20051124	JP 2004-528509	

20030808 <--				
AT 314808	T	20060215	AT 2003-787805	
20030808 <--				
ES 2250921	T3	20060416	ES 2003-787805	
20030808 <--				
NZ 537608	A	20060428	NZ 2003-537608	
20030808 <--				
RU 2316548	C2	20080210	RU 2005-106877	
20030808 <--				
ZA 2005000294	A	20060830	ZA 2005-294	
20050112 <--				
IN 2005DN00120	A	20090130	IN 2005-DN120	
20050113 <--				
MX 2005001580	A	20050425	MX 2005-1580	
20050209 <--				
US 20050234110	A1	20051020	US 2005-524345	
20050211 <--				
KR 853967	B1	20080825	KR 2005-702419	
20050212 <--				
HK 1080329	A1	20071109	HK 2006-100382	
20060110 <--				
PRIORITY APPLN. INFO.:			EP 2002-356159	A
20020812 <--			FR 2003-5233	A
20030429			WO 2003-EP9516	W
20030808				
OTHER SOURCE(S):	MARPAT 140:181332			
GI				



AB Title compds. I [wherein X = independently halo, halogeno/alkyl; Y = independently halo, halogeno/alkyl, alkoxy, phenoxy, alkylthio, dialkylamino, acyl, CN, NO<sub>2</sub>, alkylsulfonyl, phenylsulfonyl, benzylsulfonyl, S-Ph substituted by a halogen; p = 1-4; q = 1-5; with the exception of N-[[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl]-2,6-dichlorobenzamide] were prepared as fungicides, in particular as fungicidal compns. for controlling phytopathogenic fungi of crops. For example, II was prepared in 4

steps by reaction of 2,3-dichloro-5-(trifluoromethyl)pyridine with Me cyanoacetate in DMF, decarboxylation in H<sub>2</sub>O/DMSO, Pd/C hydrogenation, and acylation with 2-chlorobenzoyl chloride. In vivo tests of activity upon *Alternaria brassicae*, *Botrytis cinerea*, *Pyrenophora teres*, and *Septoria nodorum* by selected I are reported, demonstrating their fungicide efficiency (no data). Fungicidal compns. contain 0.05 to 99% active pyridylethylbenzamide.

REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS

RE FORMAT

RECORD. ALL CITATIONS AVAILABLE IN THE

L4 ANSWER 3 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 2004:80450 CAPLUS Full-text  
 DOCUMENT NUMBER: 140:145835  
 TITLE: Preparation of dibenzofused

modulators of

bicyclo[2.2.2]octane-derived amides as

the glucocorticoid receptor

INVENTOR(S): Vaccaro, Wayne; Yang, Bingwei Vera; Kim, Soong-hoon;

Huynh, Tram; Tortolani, David R.; Leavitt,

Kenneth J.;

Li, Wenying; Doweiko, Arthur M.; Chen, Xiao-

tao;

Doweiko, Lidia

PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA; et al.

SOURCE: PCT Int. Appl., 265 pp.

CODEN: PIXXD2

DOCUMENT TYPE:

Patent

LANGUAGE:

English



FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004009017	A2	20040129	WO 2003-US22300	
20030717 <--				
WO 2004009017	A3	20040708		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI,				



SK, TR,  
 BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN,  
 TD, TG  
 AU 2003251970 A1 20040209 AU 2003-251970  
 20030717 <--  
 US 20040132758 A1 20040708 US 2003-621909  
 20030717 <--  
 US 6995181 B2 20060207  
 EP 1534273 A2 20050601 EP 2003-765638  
 20030717 <--  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,  
 MC, PT,  
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK  
 JP 2006508042 T 20060309 JP 2004-523482  
 20030717 <--  
 NO 2005000074 A 20050309 NO 2005-74  
 20050106 <--  
 US 20050171136 A1 20050804 US 2005-85347  
 20050321 <--  
 PRIORITY APPLN. INFO.: US 2002-396877P P  
 20020718 <--  
 US 2003-621909 A1  
 20030717  
 WO 2003-US22300 W  
 20030717  
 OTHER SOURCE(S): MARPAT 140:145835  
 GI



AB Title compds. I [R-R4 = H, alk(en/yn)yl, alkoxy, aryl, etc.; Z = carboxamido, alkylamino, etc.] are prepared For instance, 2-amino-4,5-dimethylthiazole is coupled to the acid derived from the cycloaddn. of methacrylic acid and anthracene (CH3CN, EDCI, Et3N, HOAt, 18 h) to give II. I are glucocorticoid receptor modulators which are useful in treating diseases requiring glucocorticoid receptor agonist or antagonist therapy such as obesity, diabetes, inflammatory and immune disorders.

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS  
 RECORD. ALL CITATIONS AVAILABLE IN THE  
 RE FORMAT

L4 ANSWER 4 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 2003:261800 CAPLUS [Full-text](#)  
 DOCUMENT NUMBER: 138:271704

TITLE: Preparation of acid amide derivatives as pesticides  
 INVENTOR(S): Nakamura, Yuji; Morita, Masayuki; Yoneda, Tetsuo;  
 Izakura, Kenji  
 PATENT ASSIGNEE(S): Ishihara Sangyo Kaisha, Ltd., Japan  
 SOURCE: PCT Int. Appl., 233 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	----
WO 2003027059	A1	20030403	WO 2002-JP9560	
20020918 <--				
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
JP 2003176258	A	20030624	JP 2002-270729	
20020917 <--				
CA 2460789	A1	20030403	CA 2002-2460789	
20020918 <--				
AU 2002332181	A1	20030407	AU 2002-332181	
20020918 <--				
EP 1428817	A1	20040616	EP 2002-767967	
20020918 <--				
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
US 20040254237	A1	20041216	US 2004-489778	
20040317 <--				
PRIORITY APPLN. INFO.:			JP 2001-283969	A
20010918 <--			WO 2002-JP9560	W
20020918 <--				
OTHER SOURCE(S):	MARPAT 138:271704			
AB	Acid amide derivs. represented by the formula A-CO-CR1R2-NR3-CO-B [wherein A = Ph, benzyl, naphthyl, heterocyclic group, or fused heterocyclic group each optionally substituted by X, indanyl			

(which may be substituted by halogen, alkyl, or alkoxy), or tetrahydronaphthyl (which may be substituted by halogen, alkyl, or alkoxy); B = alkyl, cycloalkyl, Ph optionally substituted by Y, a heterocyclic group optionally substituted by Y, or a fused heterocyclic group optionally substituted by Y; X = halo, alkyl, haloalkyl, alkenyl, haloalkenyl, alkynyl, haloalkynyl, alkoxy, haloalkoxy, alkoxyalkoxy, haloalkoxyalkoxy, alkoxyhaloalkoxy, etc.; Y = halo, alkyl, haloalkyl, alkoxy, haloalkoxy, alkylthio, haloalkylthio, alkylsulfanyl, haloalkylsulfanyl, alkylsulfonyl, haloalkylsulfonyl, dialkylaminosulfonyl, NO<sub>2</sub>, cyano, etc.; R<sub>1</sub>, R<sub>2</sub> = alkyl, cyano, or CO<sub>2</sub>R<sub>14</sub>, provided that R<sub>1</sub> and R<sub>2</sub> in combination may form a 3- to 6-membered saturated carbon ring; R<sub>3</sub> = H, alkyl, alkoxyalkyl, alkylthioalkyl, COR<sub>15</sub>, S(O)mR<sub>16</sub>, or S(O)nNR<sub>17</sub>/R<sub>18</sub>; wherein R<sub>14</sub> = H, alkyl; R<sub>15</sub> = H, alkyl, alkoxy; R<sub>16</sub>, R<sub>17</sub>, R<sub>18</sub> = alkyl, haloalkyl, optionally substituted Ph or salts thereof are prepared. These compds. including N-phenacylbenzamides, N-phenacylnaphthalenecarboxamides, N-phenacylthiophenecarboxamides, N-phenacylpyrazinecarboxamides, N-phenacylquinolinecarboxamides, N-phenacylindolecarboxamides, N-phenacylfurancarboxamides, N-phenacylbenzofurancarboxamides, N-phenacylbenzodioxanecarboxamides, N-(naphthylcarbonylmethyl)benzamide, N-(thienylcarbonylmethyl)benzamides, N-(thienylcarbonylmethyl)pyridinecarboxamides, N-(pyridylcarbonylmethyl)benzamides, N-(benzodioxanylcabonylmethyl)benzamides, and N-(furylcabonylmethyl)benzamides are useful as active ingredients for pest control agents such as insecticides, acaricides, nematocides, and animal parasiticides. Thus, 0.11 g 2-fluorobenzoyl chloride was added dropwise to a mixture of 0.20 g 6-(2,2,3,3-tetrafluoro-5-methyl-1,4-benzodioxan-6-yl) 2-amino-2-Pr ketone, 0.10 g Et<sub>3</sub>N, and 7 mL THF and stirred at room temperature for 2 h to give 2-fluoro-N-[2-[(2,2,3,3-tetrafluoro-5-methyl-1,4-benzodioxan-6-yl)carbonyl]-2-propyl]benzamide (II). II at 1,600 ppm (soil application) completely controlled nematode in tomato seedlings.

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE  
FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE  
RE FORMAT



L4 ANSWER 5 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 2001:380581 CAPLUS Full-text  
DOCUMENT NUMBER: 135:5611  
TITLE: Preparation of pesticidal  
aminoheterocyclamides  
INVENTOR(S): Ducray, Pierre; Bouvier, Jacques; Mueller, Urs  
PATENT ASSIGNEE(S): Novartis A.-G., Switz.; Novartis-Erfindungen  
Verwaltungsgesellschaft m.b.H.  
SOURCE: PCT Int. Appl., 66 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
------------	------	------	-----------------	------

```

-----
WO 2001036415      A1      20010525      WO 2000-EP11387
20001116 <---
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN,
      CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH,
GM, HR,
      HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT,
      LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT,
RO, RU,
      SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US,
UZ, VN,
      YU, ZA, ZW
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE,
CH, CY,
      DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE,
TR, BF,
      BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
CA 2386318      A1      20010525      CA 2000-2386318
20001116 <---
BR 2000015671      A      20020723      BR 2000-15671
20001116 <---
EP 1230239      A1      20020814      EP 2000-983143
20001116 <---
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,
MC, PT,
      IE, SI, LT, LV, FI, RO, MK, CY, AL
JP 2003514816      T      20030422      JP 2001-538904
20001116 <---
AU 764228      B2      20030814      AU 2001-20005
20001116 <---
NZ 518376      A      20040130      NZ 2000-518376
20001116 <---
RU 2259370      C2      20050827      RU 2002-116255
20001116 <---
ZA 2002003861      A      20021205      ZA 2002-3861
20020515 <---
US 6667326      B1      20031223      US 2002-130492
20020516 <---
MX 2002005023      A      20020918      MX 2002-5023
20020517 <---
PRIORITY APPLN. INFO.:      CH 1999-2107      A
19991118 <---
WO 2000-EP11387      W
20001116 <---
OTHER SOURCE(S):      MARPAT 135:5611
GI

```



AB The title compds. [I; H, halo, alkyl, etc.; R2 = H, alkyl, alkylenophenyl, etc.; X1 = N, C(CN); X2 = N, C(CN), C(CO2R6), etc.; X3 = O, S; q = CH, N; R3, R4 = H, alkyl; or R3R4 together with the C-atom to which they are bonded = cycloalkyl; R5 = alkyl, alkenyl, alkynyl, etc.; R6 = alkyl, Ph, CH2Ph; m = 1-3; n = 0-1] which have advantageous pesticidal properties and are especially suitable for the control of pests on domestic and farm animals, were prepared and formulated. Thus, treating 1-(3,5-dichloropyrid-2-yl)cyclopropyl-1-carboxylic acid with (COCl)2 and a drop of DMF followed by reacting the resulting intermediate with 2-amino-5-cyano-4-trifluoromethylthiazole in the presence of diisopropylethylamine and 4-dimethylaminopyridine in CH2Cl2 afforded II. Biol. data for compds. I were given.

REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

L4 ANSWER 6 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2001:136943 CAPLUS [Full-text](#)

DOCUMENT NUMBER: 134:174246

TITLE: Preparation of pyridine derivative fungicides

INVENTOR(S): Cooke, Tracey; Hardy, David; Moloney, Brian; Thomas,

Peter Stanley; Steele, Chris Richard; Briggs,

Geoffrey

Gower

PATENT ASSIGNEE(S): Aventis CropScience GmbH, Germany

SOURCE: PCT Int. Appl., 56 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	----
WO 2001011965	A1	20010222	WO 2000-EP8143	

20000809 <--  
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW  
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG  
BR 2000013371 A 20020507 BR 2000-13371  
20000809 <--  
EP 1204323 A1 20020515 EP 2000-960499  
20000809 <--  
EP 1204323 B1 20040714  
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL  
JP 2003506465 T 20030218 JP 2001-516328  
20000809 <--  
AT 270817 T 20040715 AT 2000-960499  
20000809 <--  
PT 1204323 T 20041130 PT 2000-960499  
20000809 <--  
ES 2220533 T3 20041216 ES 2000-960499  
20000809 <--  
CN 1209016 C 20050706 CN 2000-811802  
20000809 <--  
IN 2002MN00092 A 20050318 IN 2002-MN92  
20020125 <--  
MX 2002001453 A 20030128 MX 2002-1453  
20020211 <--  
US 6821992 B1 20041123 US 2002-49976  
20020709 <--  
PRIORITY APPLN. INFO.: GB 1999-19499 A  
19990818 <--  
GB 1999-19500 A  
19990818 <--  
WO 2000-EP8143 W  
20000809 <--  
OTHER SOURCE(S): MARPAT 134:174246  
AB The pyridine derivs. A1CR1R2LA2 (A1 = (un)substituted 2-pyridyl or its N-oxide; Y = LA2 or L1A3; A2, A3 = (un)substituted carbocyclyl or heterocyclyl; L = NR5C(:X)NR6, NR5C(:X)CHR3, CHR3NR5CHR4, etc.; L1 = NR9C(:X)X1CHR7, NR9C(:X)CHR7CHR8, etc.; R1-9 = CN, NO2, halo, etc.) are prepared as agrochem. fungicides.  
REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 7 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2000:790308 CAPLUS Full-text

DOCUMENT NUMBER: 133:350214

TITLE: Preparation of  
4-benzimidazolylmethoxy-3-

halophenylmethoxybenzoates

and analogs as tRNA synthetase inhibitors  
INVENTOR(S): Leeman, Aaron H.; Hammond, Milton L.; Maletic,  
Milana;

Santorelli, Gina M.; Waddell, Sherman F.;

Finn, John;

PATENT ASSIGNEE(S): Morytko, Michael; Hill, Jason; Keith, Dennis  
Merck & Co., Inc., USA; Cubist Pharmaceuticals  
Inc.

SOURCE: PCT Int. Appl., 90 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent


LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
WO 2000066120	A1	20001109	WO 2000-US12178	
20000505 <--				
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH,				
CN, CR,				
CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,				
HR, HU,				
ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT,				
LU, LV,				
MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD,				
SE, SG,				
SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU,				
ZA, ZW				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH,				
CY, DE,				
DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF,				
BJ, CF,				
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
CA 2372079	A1	20001109	CA 2000-2372079	
20000505 <--				
EP 1176958	A1	20020206	EP 2000-930366	
20000505 <--				
EP 1176958	B1	20040728		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,				
MC, PT,				
IE, SI, LT, LV, FI, RO				
US 6348482	B1	20020219	US 2000-566275	
20000505 <--				
JP 2002543130	T	20021217	JP 2000-615005	
20000505 <--				
AT 271868	T	20040815	AT 2000-930366	
20000505 <--				
AU 776773	B2	20040923	AU 2000-48199	
20000505 <--				

EP 1466603 A2 20041013 EP 2004-76350  
 20000505 <--  
 EP 1466603 A3 20041020  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,  
 MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL  
 ES 2226839 T3 20050401 ES 2000-930366  
 20000505 <--  
 US 20020040147 A1 20020404 US 2001-934743  
 20010822 <--  
 US 6545015 B2 20030408  
 PRIORITY APPLN. INFO.: US 1999-132545P P  
 19990505 <--  
 EP 2000-930366 A3  
 20000505 <--  
 US 2000-566275 A3  
 20000505 <--  
 WO 2000-US12178 W  
 20000505 <--  
 OTHER SOURCE(S): MARPAT 133:350214  
 GI



AB RCR5R6ZOOCR7R8R9 [I; R = (hetero)aryl; R5-R8 = H or alkyl; R9 = (un)substituted CH2NHC(:NH)NH2, N-containing heteroaryl(aminomethyl), etc.; Z = (un)substituted 1,2-phenylene] were prepared as bactericides and fungicides. Thus, 3,4-(HO)(MeOCH2O)C6H3CO2Et was O-alkylated by 2,4-Cl2C6H3CH2Cl and the O-deprotected product O-alkylated by 2-chloromethyl-1-[(2-trimethylsilylethoxy)methyl]benzimidazole (preparation given) to give, after deprotection and saponification, title compound II. Data for biol. activity of I were given.

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

L4 ANSWER 8 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1964:23151 CAPLUS Full-text

DOCUMENT NUMBER: 60:23151

ORIGINAL REFERENCE NO.: 60:4056e-g

TITLE: Substituted salicylamides and their analgesic effect

AUTHOR(S): Profft, E.; Hoegel, E.

SOURCE: Pharmazie (1962), 17(12), 731-4

CODEN: PHARAT; ISSN: 0031-7144

DOCUMENT TYPE: Journal



LANGUAGE: Unavailable

AB Salicyloyl chloride (I) (0.1 mole) dissolved in 25-100 ml. ether was added over 15-45 min. to 0.2 mole amine in 175-400 ml. ether at 10-15°, and the mixture stirred 1-2 hrs. (Method A). I (1 mole) was added to a mixture of 1 mole amine in ether containing the calculated amount of 20% aqueous NaOH at -10 to -15° with stirring (Method B). Thus were prepared the following salicylamides (amine used, method, % yield, and m. p. given): piperidine, A, 96, 143.5-4.5°; hexamethylenimine, A, 96, 117.5-18.5°; 2-pyridylmethylamine, A, 85, 115-15.5°; o-anisidine, A, 77, 112-13°; p-anisidine, A, 89, 160-60.5°; 4-aminophenyl Et ether, B, 95, 141-2°; 4-aminophenyl Pr ether, B, 88, 135-6°; 2-aminophenyl Bu ether, A, 96, 112.5-13°; 4-aminophenyl Bu ether, B, 94, 133-3.5°; 4-butoxybenzylamine, A, 86, 86-7°; 2-pyridylethylbenzylamine, A, 43, 106-7°. p-Alkoxyanilides showed better analgesic activity (hot-plate method) than three similar com. compds. Heterocyclic salicylamides and, particularly, salicylopiperidide showed good effects.

L4 ANSWER 9 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1960:7335 CAPLUS Full-text  
DOCUMENT NUMBER: 54:7335  
ORIGINAL REFERENCE NO.: 54:1556d-g  
TITLE: Pyridylethylated salicylamides  
INVENTOR(S): Shapiro, Seymour L.; Freedman, Louis; Rose, Ira M.  
PATENT ASSIGNEE(S): U.S. Vitamin & Pharmaceutical Corp.  
SOURCE: Continuation-in-part of U.S. 2,835,668 (C.A. 53,  
2261b)  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	----
-----				
US 2899437		19590811	US	
00000000 <--				

AB The title compds. are useful as analgesics, fungicidess, anesthetics, and for central nervous system depression, ganglionic blockade, and antiinflammatory response. Thus, 0.005 mole of an N-pyridylethyl-1,3-benzoxazine-2,4-dione in NaOH solution is stirred until solution occurs, (2-48 hrs.), acidified with HCl, and filtered to yield the desired N-pyridylsalicylamide (I). I with MeI gives the corresponding N-methylpyridinium iodide. The new compds. prepared are (bz-substituent, pyridine group, % yield, and m.p. given): 2-C5H4N (II), 5-Cl, 87, 139°; 5-Cl, 2-pyridyl-5-ethyl (III), 81, 131°; 5-Cl, 4-C5H4N (IV), 70, 154-5° (methiodide m. 196-201°); 5-Br, II, 51, 143°; 5-Br, III, 67, 131° (methiodide m. 198-201°); 3-Me, II, 73, 97° (methiodide m. 188°); 3-Me, III, 68, 106° (methiodide m. 198-200°); 3-Me, IV, 65, 152-3°; 5-Ph, III, 70, 147-8°; 5-Ph, IV, 66, 127-8°; 4-OH, II, 60, 209-10°; 4-OH, III, -, 172-3°; 4-OH, IV, 71, 275-6°; 5-OH, II, 53, 202-5°; 5-OH, III, 55, 160-1°; 5-OH, IV, 58, 238-42°. Cf. C.A. 51, 14731b.

```
=> s l2 and ('methionin?')
      444 L2
      61 'METHIONIN?'
L5      0 L2 AND ('METHIONIN?')


=> s l2 and methionine
      444 L2
      97295 METHIONINE
      557 METHIONINES
      97489 METHIONINE
      (METHIONINE OR METHIONINES)
L6      3 L2 AND METHIONINE
```

```
=> d l6 ibib abs
```

```
L6 ANSWER 1 OF 3 CAPLUS COPYRIGHT 2009 ACS on STN
ACCESSION NUMBER: 2005:891135 CAPLUS Full-text
DOCUMENT NUMBER: 143:207621
TITLE: Synergistic fungicidal compositions comprising
a      pyridylethylbenzamide derivative and a
      methionine biosynthesis inhibitor
INVENTOR(S): Gouot, Jean-Marie; Grosjean-Cournoyer, Marie-
Claire
PATENT ASSIGNEE(S): Bayer Cropscience SA, Fr.
SOURCE: PCT Int. Appl., 19 pp.
      CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:
```

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	----
WO 2005077182	A1	20050825	WO 2005-EP2567	
20050210				
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,			
CA, CH,	CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,			
GB, GD,	GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,			
KZ, LC,	LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,			
NA, NI,	NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK,			
SL, SY,	TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA,			
ZM, ZW	RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,			
ZW, AM,	AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ,			
DE, DK,				

EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL,  
 PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
 GW, ML, MR, NE, SN, TD, TG  
 EP 1570737 A1 20050907 EP 2004-356015  
 20040212 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,  
 MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK  
 AU 2005213185 A1 20050825 AU 2005-213185  
 20050210 CA 2551148 A1 20050825 CA 2005-2551148  
 20050210 EP 1713335 A1 20061025 EP 2005-715941  
 20050210 EP 1713335 B1 20080827  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,  
 MC, PT, IE, SI, LT, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK, IS  
 CN 1917761 A 20070221 CN 2005-80004217  
 20050210 BR 2005006619 A 20070502 BR 2005-6619  
 20050210 JP 2007522186 T 20070809 JP 2006-552586  
 20050210 AT 406097 T 20080915 AT 2005-715941  
 20050210 IN 2006DN03605 A 20070831 IN 2006-DN3605  
 20060622 MX 2006009066 A 20061113 MX 2006-9066  
 20060809 KR 2006126579 A 20061207 KR 2006-717764  
 20060901 KR 838539 B1 20080617  
 US 20070137273 A1 20070621 US 2006-588360  
 20061010 PRIORITY APPLN. INFO.: EP 2004-356015 A  
 20040212 US 2004-636999P P  
 20041217 WO 2005-EP2567 W  
 20050210  
 OTHER SOURCE(S): MARPAT 143:207621  
 GI



AB Synergistic fungicidal compns. comprise a pyridylethylbenzamide derivative I (X = halo, alkyl or haloalkyl; Y = X, alkenyl, alkynyl, alkoxy, etc.; p = 1-4; q = 1-5) and a compound capable of inhibiting methionine biosynthesis. Optionally, the composition further comprises an addnl. fungicide.

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS


RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> d 16 ibib abs 1-3

L6 ANSWER 1 OF 3 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 2005:891135 CAPLUS Full-text  
 DOCUMENT NUMBER: 143:207621  
 TITLE: Synergistic fungicidal compositions comprising  
 a pyridylethylbenzamide derivative and a methionine biosynthesis inhibitor  
 Gouot, Jean-Marie; Grosjean-Cournoyer, Marie-  
 INVENTOR(S): Claire  
 PATENT ASSIGNEE(S): Bayer Cropscience SA, Fr.  
 SOURCE: PCT Int. Appl., 19 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005077182	A1	20050825	WO 2005-EP2567	
20050210				
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
EP 1570737	A1	20050907	EP 2004-356015	

20040212  
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,  
MC, PT,  
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK  
AU 2005213185 A1 20050825 AU 2005-213185  
20050210  
CA 2551148 A1 20050825 CA 2005-2551148  
20050210  
EP 1713335 A1 20061025 EP 2005-715941  
20050210  
EP 1713335 B1 20080827  
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,  
MC, PT,  
IE, SI, LT, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK, IS  
CN 1917761 A 20070221 CN 2005-80004217  
20050210  
BR 2005006619 A 20070502 BR 2005-6619  
20050210  
JP 2007522186 T 20070809 JP 2006-552586  
20050210  
AT 406097 T 20080915 AT 2005-715941  
20050210  
IN 2006DN03605 A 20070831 IN 2006-DN3605  
20060622  
MX 2006009066 A 20061113 MX 2006-9066  
20060809  
KR 2006126579 A 20061207 KR 2006-717764  
20060901  
KR 838539 B1 20080617  
US 20070137273 A1 20070621 US 2006-588360  
20061010  
PRIORITY APPLN. INFO.: EP 2004-356015 A  
20040212 US 2004-636999P P  
20041217 WO 2005-EP2567 W  
20050210  
OTHER SOURCE(S): MARPAT 143:207621  
GI



AB Synergistic fungicidal compns. comprise a pyridylethylbenzamide derivative I (X = halo, alkyl or haloalkyl; Y = X, alkenyl, alkynyl, alkoxy, etc.; p = 1-4; q = 1-5) and a compound capable of inhibiting methionine biosynthesis. Optionally, the composition further comprises an addnl. fungicide.

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS

## RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

L6 ANSWER 2 OF 3 CAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 2004:465510 CAPLUS Full-text  
DOCUMENT NUMBER: 141:133551  
TITLE: Thyroid receptor ligands. Part 2:  
thyromimetics with improved selectivity for the thyroid hormone  
receptor  
beta  
AUTHOR(S): Hangeland, Jon J.; Doweiko, Arthur M.;  
Dejneka, Tamara; Friends, Todd J.; Devasthale, Pratik;  
Mellstrom, Karin; Sandberg, Johnny; Grynfarb,  
Marlena; Sack, John S.; Einspahr, Howard; Faernegardh,  
Mathias; Husman, Bolette; Ljunggren, Jan; Koehler,  
Konrad; Sheppard, Cheryl; Malm, Johan; Ryono, Denis E.  
CORPORATE SOURCE: Pharmaceutical Research Institute, Bristol-  
Myers Squibb, Princeton, NJ, 08543, USA  
SOURCE: Bioorganic & Medicinal Chemistry Letters  
(2004), 14(13), 3549-3553  
CODEN: BMCLE8; ISSN: 0960-894X  
PUBLISHER: Elsevier Science B.V.  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
OTHER SOURCE(S): CASREACT 141:133551  
AB A set of thyromimetics having improved selectivity for TR- $\beta$ 1 were prepared by replacing the 3'-iso-Pr group of 2 and 3 with substituents having increased steric bulk. From this limited SAR study, the most potent and selective compds. identified were derived from 2 and contained a 3'-Ph moiety bearing small hydrophobic groups meta to the biphenyl link. X-ray crystal data of 15c complexed with TR- $\beta$ 1 LBD shows methionine 442 to be displaced by the bulky R3' Ph Et amide side chain. Movement of this amino acid side chain provides an expanded pocket for the bulky side chain while the ligand-receptor complex retains full agonist activity.

REFERENCE COUNT: 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS

## RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

L6 ANSWER 3 OF 3 CAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 1999:529132 CAPLUS Full-text  
DOCUMENT NUMBER: 131:170355  
TITLE: Preparation of heterocycle-containing  
benzamide derivatives as farnesyl transferase inhibitors  
INVENTOR(S): Drake, David John; Wardleworth, James Michael  
PATENT ASSIGNEE(S): Zeneca Limited, UK; Zeneca Pharma S.A.

SOURCE: PCT Int. Appl., 138 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9941235	A1	19990819	WO 1999-GB369	
19990204				
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
AU 9924351	A	19990830	AU 1999-24351	
19990204				
EP 1054865	A1	20001129	EP 1999-903834	
19990204				
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
JP 2002503650	T	20020205	JP 2000-531430	
19990204				
ZA 9901032	A	19990810	ZA 1999-1032	
19990209				
PRIORITY APPLN. INFO.:			EP 1998-400294	A
19980210			WO 1999-GB369	W
19990204				
OTHER SOURCE(S):	MARPAT 131:170355			
GI				

\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

AB The present invention relates to compds. of formula (I; wherein A is of formula Q, Q1, or Ar1CH2E(Ar2); B is Ph, pyridyl, pyridazinyl, pyrimidyl, pyrazinyl, thienyl, thiazolyl, furyl or oxazolyl, the ring being substituted on ring carbon atoms by R1 and -(CH2)nR2; or B is pyrrolyl, pyrazolyl or imidazolyl, and when A is of formula Q or Q1, B can also be naphthyl substituted by R1 and (CH2)nR2; R1 is of the formula -CONHCH(R10)R11; ; R2 is Ph or

heteroaryl; n is 0, 1 or 2; wherein R3 is hydrogen, C2-5 alkanoyl, C1-4 alkoxy carbonyl, C2-4 alkenyloxy carbonyl, phenyl-C1-3 alkyl, phenoxy carbonyl, etc.; R4 is hydrogen, C1-4 alkyl, C2-5 alkanoyl, C1-4 alkoxy carbonyl, phenyl-C1-3 alkyl, benzoyl, heteroaryl C1-3 alkyl or heteroaryl; D is a linking moiety selected from (un)substituted Q3 - Q5; Ar1 is (un)substituted imidazol-1-, -2-, or -3-yl; Ar2 is Ph or heteroaryl; E is C:CH, CHCH2, N-(un)substituted CHNH or CHNHCH2, CHO, CHOCH2; wherein R10 is hydrogen or (CH2)qR12 (q is 0-4) and R11 is of the formula CH2OR13, COR14, CH2COR14, is morpholino-C1-4 alkyl, pyrrolidin-1-yl-C1-4 alkyl, piperidin-1-yl-C1-4 alkyl, etc.; R12 is hydrogen, C1-4 alkylsulfanyl, C1-4 alkyl sulfonyl, hydroxy, C1-4 alkoxy, etc.; R13 is hydrogen, C1-4 alkyl, Ph, heteroaryl, C2-5 alkanoyl, etc.; R14 (un)substituted C1-4 alkyl, Ph, phenyl-C1-3 alkyl, cyano, C2-4 alkanoyloxy, HO, etc.) or pharmaceutically acceptable salts or prodrugs thereof. These compds. are useful for the treatment of a disease mediated through farnesylation of mutant ras products by inhibition of the enzyme farnesyl-protein transferase (FPTase), especially cancer. Thus, 4-([1-(4-Fluorophenyl)-2-(imidazol-1-yl)ethylamino]-2-(4-fluorophenyl)benzoic acid was condensed with L-methionine Me ester hydrochloride using 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide, HOBt, and 4-dimethylaminopyridine in CH2Cl2 at ambient temperature for 5 h to give 80% N-(4-([1-(4-Fluorophenyl)-2-(imidazol-1-yl)ethylamino]-2-(4-fluorophenyl)benzoyl)-L-methionine Me ester which was reduced by LiBH4 in THF at 0° at ambient temperature overnight to give N-benzoyl-L-methioninol derivative (II).

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> s l2 and (methionine or amino acid or protein)

<-----User Break----->

SEARCH ENDED BY USER

=> s l2 and (methionine or amino acid or protein)

<-----User Break----->

SEARCH ENDED BY USER

=> s l2 and (amino acid? or protein?)

<-----User Break----->

SEARCH ENDED BY USER

=> s l2

L7 444 L2

=> s l2 and methionine

444 L2

97295 METHIONINE

557 METHIONINES



97489 METHIONINE  
(METHIONINE OR METHIONINES)

L8 3 L2 AND METHIONINE

=> s l2 and (protein? or 'amino acid?')  
<-----User Break----->

SEARCH ENDED BY USER


=> s l2 and (germinat? or mycelium)  
444 L2  
63685 GERMINAT?  
16266 MYCELIUM  
29 MYCELIUMS  
9113 MYCELIA  
2 MYCELIAS  
23446 MYCELIUM  
(MYCELIUM OR MYCELIUMS OR MYCELIA OR MYCELIAS)  
L9 1 L2 AND (GERMINAT? OR MYCELIUM)

=> d l9 ibib abs

L9 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 2005:891134 CAPLUS Full-text  
DOCUMENT NUMBER: 143:207620  
TITLE: Synergistic fungicidal compositions comprising  
a  
pyridylethylbenzamide derivative and a  
compound  
capable of inhibiting spore germination or  
mycelium growth by acting on different  
metabolic routes  
Grosjean-Cournoyer, Marie-Claire; Guot, Jean-  
Marie  
INVENTOR(S):  
PATENT ASSIGNEE(S): Bayer Cropscience SA, Fr.  
SOURCE: PCT Int. Appl., 35 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	----
WO 2005077181	A1	20050825	WO 2005-EP2566	
20050210				
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,			
CA, CH,	CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,			
GB, GD,	GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,			
KZ, LC,	LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,			
NA, NI,	NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK,			
SL, SY,				

TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA,  
 ZM, ZW  
 RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,  
 ZW, AM,  
 AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ,  
 DE, DK,  
 EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL,  
 PL, PT,  
 RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
 GW, ML,  
 MR, NE, SN, TD, TG  
 EP 1570738 A1 20050907 EP 2004-356017  
 20040212  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,  
 MC, PT,  
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK  
 AU 2005213184 A1 20050825 AU 2005-213184  
 20050210  
 CA 2551147 A1 20050825 CA 2005-2551147  
 20050210  
 EP 1713334 A1 20061025 EP 2005-715940  
 20050210  
 EP 1713334 B1 20080723  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,  
 MC, PT,  
 IE, SI, LT, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK, IS  
 CN 1917762 A 20070221 CN 2005-80004260  
 20050210  
 BR 2005006613 A 20070502 BR 2005-6613  
 20050210  
 JP 2007522185 T 20070809 JP 2006-552585  
 20050210  
 AT 401791 T 20080815 AT 2005-715940  
 20050210  
 ES 2311213 T3 20090201 ES 2005-715940  
 20050210  
 IN 2006DN03600 A 20070831 IN 2006-DN3600  
 20060622  
 MX 2006009067 A 20061113 MX 2006-9067  
 20060809  
 KR 838540 B1 20080617 KR 2006-716373  
 20060814  
 US 20070142444 A1 20070621 US 2006-588532  
 20061012  
 PRIORITY APPLN. INFO.: EP 2004-356017 A  
 20040212 US 2004-636898P P  
 20041218 WO 2005-EP2566 W  
 20050210  
 OTHER SOURCE(S): MARPAT 143:207620  
 GI



AB Synergistic fungicidal compns. comprise at least a pyridylethylbenzamide derivative I (X = halo, alkyl or haloalkyl; Y = X, alkenyl, alkynyl, alkoxy, etc.; p = 1-4; q = 15) and a compound capable of inhibiting spore germination or mycelium growth by acting on different metabolic routes. A composition optionally contains an addnl. fungicidal compound

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> s fungici? and (germinat? or mycelium?)  
118522 FUNGICI?  
63685 GERMINAT?  
16283 MYCELIUM?  
L10 5406 FUNGICI? AND (GERMINAT? OR MYCELIUM?)

=> s l10 and (methionine)  
97295 METHIONINE  
557 METHIONINES  
97489 METHIONINE  
(METHIONINE OR METHIONINES)  
L11 18 L10 AND (METHIONINE)

=> d l11 ibib abs 1-18

L11 ANSWER 1 OF 18 CAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 2006:650551 CAPLUS Full-text  
DOCUMENT NUMBER: 146:289803  
TITLE: Characterization of laboratory mutants of Botrytis cinerea resistant to QoI fungicides  
AUTHOR(S): Markoglou, Anastasios N.; Malandrakis, Vitoratos, Andreas G.; Ziogas, Basil N.  
CORPORATE SOURCE: Laboratory of Pesticide Science, Agricultural University of Athens, Athens, 118 55, Greece  
SOURCE: European Journal of Plant Pathology (2006), 115(2), 149-162  
CODEN: EPLPEH; ISSN: 0929-1873  
PUBLISHER: Springer  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Botrytis cinerea mutants with moderate and high resistance to pyraclostrobin, a Qo inhibitor of mitochondrial electron transport at the cytochrome bcl complex, were isolated at high frequency

after nitrosoguanidine-induced mutagenesis and selection on medium containing pyraclostrobin and salicylhydroxamate (SHAM), a specific inhibitor of cyanide-resistant (alternative) respiration. Oxygen uptake in whole fungal cells was strongly inhibited in the wild-type strain by pyraclostrobin and SHAM, but not in the mutant isolates. Cross-resistance studies with other Qo and Qi inhibitors (QoI and QiI) of cytochrome bcl complex of mitochondrial respiration showed that the mutation(s) for resistance to pyraclostrobin also decreased the sensitivity of the mutant strains to other QoI (azoxystrobin, fluoxastrobin, trifloxystrobin, picoxystrobin), but not to famoxadone and to the QiI cyazofamid and antimycin-A. Increased sensitivity of pyraclostrobin-resistant strains to the carboxamide boscalid (inhibitor of complex II) and to the anilinopyrimidine cyprodinil (methionine biosynthesis inhibitor) was observed. No effect of pyraclostrobin resistance mutation(s) on fungicidal activity of the hydroxyanilide fenhexamid, the phenylpyrrole fludioxonil, the benzimidazole benomyl, and the phenylpyridinamine fluazinam, which affect other cellular pathways, was observed. Study of fitness parameters in the wild-type and pyraclostrobin-resistant mutants of *B. cinerea* showed that most mutants had decreased sporulation, conidial germination, and sclerotia production. Stability studies of the pyraclostrobin-resistant phenotype showed decreased resistance, mainly in moderate resistant strains, when the mutants were grown on inhibitor-free media. A rapid recovery of the resistance level was observed after the mutants were returned to selective media. Study of competitive ability of mutant isolates against the wild-type parent strain (use of mixed conidial population) showed that all mutants were less competitive than the wild-type strain in vitro. The competitive ability of highly resistant mutants was higher than in moderate mutants. Pathogenicity tests on cucumber seedlings showed that all mutant strains had an infection ability similar to the wild-type parent strain. Preventive applications of the com. product of F-500 25EC (pyraclostrobin) were effective against lesion development on cotyledons by the wild-type, but ineffective, even at high concns., against disease caused by the pyraclostrobin-resistant isolates. Boscalid (F-510 50WG) was equally effective against the disease caused by the wild-type or pyraclostrobin-resistant mutants.

REFERENCE COUNT: 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 2 OF 18 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 2005:1322975 CAPLUS [Full-text](#)  
 DOCUMENT NUMBER: 144:186384  
 TITLE: Antifungal effects of cysteine towards *Eutypa* lata, a pathogen of vineyards  
 AUTHOR(S): Octave, Stephane; Amborabe, Benigne-Ernest; Luini, Estelle; Ferreira, Thierry; Fleurat-Lessard, Pierrette; Roblin, Gabriel  
 CORPORATE SOURCE: Laboratoire de Physiologie, Biochimie, Biologie

levures,  
Poitiers,  
SOURCE:  
PUBLISHER:  
DOCUMENT TYPE:  
LANGUAGE:  
AB

Moleculaire Vegetales et Genetique des  
Universite de Poitiers (CNRS, UMR 6161),  
86022, Fr.  
Plant Physiology and Biochemistry (Amsterdam,  
Netherlands) (2005), 43(10-11), 1006-1013  
CODEN: PPBIEX; ISSN: 0981-9428  
Elsevier B.V.  
Journal  
English

Cysteine inhibited mycelial growth of the pathogenic fungus affecting grapevines *E. lata* in a concentration-dependent manner. The threshold value (defined by the concentration inducing a growth inhibition >5%) was 0.5 mM. A 10 mM concentration induced a complete inhibition of growth and triggered necrotic processes as evidenced by an increasing number of nuclei stained by propidium iodide. In conditions mimicking the plant environment (in particular, a pH near the apoplastic value, i.e. 5.5), 6 mM cysteine induced dramatic modifications in the structural organization of the mycelium (wall, mitochondria, vacuoles, and nucleus) leading to death of the hyphae. The antifungal effect of the mol. increased at the acidic exptl. pH (pH 4.1). The effect was highly specific to cysteine since modifying the mol. arrangement or masking the SH-function hindered the antifungal efficiency. Cysteine spectrum of action was broad among the various strains of *E. lata* tested. However, a lower efficiency was observed against fungal species intervening in other grapevine diseases (esca, black dead arm). Besides its direct antifungal effect, the role of cysteine presents particular interest in the fight against fungal pathogens since it triggered an excretion of ergosterol, a compound with elicitor properties. Therefore, cysteine may indirectly increase plant defense reactions.

REFERENCE COUNT: 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE  
RE FORMAT

L11 ANSWER 3 OF 18 CAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 2005:25408 CAPLUS Full-text  
DOCUMENT NUMBER: 142:426689  
TITLE: Aviglycine and propargylglycine inhibit conidial germination and mycelial growth of *Fusarium oxysporum* f. sp. *luffae*  
AUTHOR(S): Jin, Jung-Kang; Adams, Douglas O.; Ko, Yeong; Yu, Chih-Wen; Lin, Chin-Ho  
CORPORATE SOURCE: Department of Life Science, National Chung Hsing University, Taichung, Taiwan  
SOURCE: Mycopathologia (2004), 158(3), 369-375  
CODEN: MYCPAH; ISSN: 0301-486X  
PUBLISHER: Kluwer Academic Publishers  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Two inhibitors, aviglycine and propargylglycine, were tested for their ability to suppress methionine synthesis thus inhibit conidial germination and mycelial growth of Czapek-Dox liquid medium grown *Fusarium oxysporum* f. sp. *luffae*. Aviglycine inhibited conidial germination in the range of 0.3-7.6  $\mu$ M. The linear inhibition range for mycelial growth was about 7.6-762.9  $\mu$ M. Although aviglycine did not completely inhibit both conidial germination and mycelial growth, it showed significant inhibitory effect at 1.5  $\mu$ M. The inhibition range for propargylglycine against conidial germination and mycelial growth were from 0.08 to 8841  $\mu$ M and from 0.8 to 884.1  $\mu$ M, resp. Propargylglycine inhibited conidial germination and mycelial growth at a concentration of 8841  $\mu$ M. The EC50 values of aviglycine were 1  $\mu$ M for conidial growth and 122  $\mu$ M for mycelial growth, and the EC50 values of propargylglycine were 47.7  $\mu$ M for conidial growth and 55.6  $\mu$ M for mycelial growth. Supplement of methionine released inhibition of aviglycine or propargylglycine to conidial germination. In addition, a mixture of aviglycine (1.5  $\mu$ M) and propargylglycine (8841  $\mu$ M) showed additive inhibitive effect than applied alone on 10 isolates. From these results, both aviglycine and propargylglycine exhibited inhibitory activity, and suggest that they can provide potential tools to design novel fungicide against fungal pathogens.

REFERENCE COUNT: 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

L11 ANSWER 4 OF 18 CAPLUS COPYRIGHT 2009 ACS ON STN

ACCESSION NUMBER: 2004:193241 CAPLUS Full-text

DOCUMENT NUMBER: 141:36274

TITLE: Polyamine metabolism during the germination of *Sclerotinia sclerotiorum* ascospores and its relation with host infection

AUTHOR(S): Garriz, Andres; Dalmasso, Maria C.; Marina, Maria;

Rivas, Elisa I.; Ruiz, Oscar A.; Pieckenstein, Fernando L.

CORPORATE SOURCE: Instituto de Investigaciones Biotecnologicas-Instituto

de Tecnologico de Chascomus, Universidad Nacional General San Martin-Consejo Nacional de

Investigaciones Cientificas y Tecnicas, Buenos Aires, Argent.

SOURCE: New Phytologist (2004), 161(3), 847-854

CODEN: NEPHAV; ISSN: 0028-646X

PUBLISHER: Blackwell Publishing Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Polyamine biosynthesis inhibitors were used to study polyamine metabolism during the germination of *Sclerotinia sclerotiorum* ascospores, and to evaluate the potential of polyamine biosynthesis inhibition for the control of ascospore-borne diseases in plants. The effects of inhibitors on ascospore

germination, free polyamine levels, ornithine decarboxylase activity and development of disease symptoms on tobacco (Nicotiana tabacum) leaf disks inoculated with ascospores were determined  $\alpha$ -Difluoromethylornithine inhibited ornithine decarboxylase and decreased free spermidine levels, but had no effect on ascospore germination. Both, the spermidine synthase inhibitor cyclohexylamine and the S-adenosyl-methionine decarboxylase inhibitor methylglyoxal bis-[guanyl hydrazone] decreased free spermidine levels, but only the latter inhibited ascospore germination, at concns. of 5 mM or higher. Lesion development on leaf disks was reduced by cyclohexylamine and methylglyoxal bis-[guanyl hydrazone], but not by  $\alpha$ -difluoromethylornithine. In the absence of inhibitors, dormant ascospores contained higher polyamine levels than mycelium. Ascospore germination did not depend on ornithine decarboxylase activity and inhibitors of this enzyme will probably have a limited potential for the control of ascospore-borne plant diseases. On the contrary, spermidine synthase and S-adenosylmethionine decarboxylase could be more suitable targets for fungicidal action. The relative insensitivity of ascospore germination to polyamine biosynthesis inhibitors may be caused by their high polyamine content.

REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 5 OF 18 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2001:627332 CAPLUS Full-text

DOCUMENT NUMBER: 135:299904

TITLE: Nitrolin and techniques for its use on winter wheat

crops  
AUTHOR(S): Niyazmetov, U. K.; Kariev, A. U.;  
Dustmukhamedov, T.

T.  
CORPORATE SOURCE: Inst. Khim. Rastitel'nykh Veshchestv im. S. Yu.

Yunusova, AN Ruz, Uzbekistan  
SOURCE: Doklady Akademii Nauk Respubliki Uzbekistan  
(2001),

(3), 34-37  
CODEN: DARUEE; ISSN: 1019-8954  
Fan

PUBLISHER: Journal

DOCUMENT TYPE: Russian

LANGUAGE: Russian  
AB Growth regulating activity of nitrolin and its compatibility with seed treatment with the fungicide tuzal were investigated. The following parameters were used: seed germination, susceptibility to root rot, grain yield, content of starch in the grain, and content of selected amino acids in the leaves. As a result of growth processes intensification, the plants treated with nitrolin had higher rate of germination compared to control plants, lower number of diseased plants, higher grain yield and higher starch content in the grain. The decrease in root rot occurrence was also supplemented by the fungicidal action of tuzal. The composition of amino acids in treated plants did not differ from

the control, although their content was higher in leaves of nitrolin treated plants.

L11 ANSWER 6 OF 18 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1997:380005 CAPLUS Full-text

DOCUMENT NUMBER: 127:1954

ORIGINAL REFERENCE NO.: 127:463a,466a

TITLE: Bioregulatory effects of the fungicidal strobilurin kresoxim-methyl in wheat (*Triticum aestivum*)

AUTHOR(S): Grossmann, Klaus; Retzlaff, Gunter

CORPORATE SOURCE: Agricultural Res. Station, BASF, Limbergerhof, D-67114, Germany

SOURCE: Pesticide Science (1997), 50(1), 11-20

CODEN: PSSCBG; ISSN: 0031-613X

PUBLISHER: Wiley

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Apart from its fungicidal effect, the strobilurin kresoxim-Me (BAS 490 F) was found to induce physiol. and developmental alterations in wheat (*Triticum aestivum* L.) which are seen in connection with improved yield. In a series of biotests including heterotrophic maize and photoautotrophic algal cell suspensions, duckweed, isolated mustard shoots and germinating cress seeds, kresoxim-Me showed a similar response pattern to standard auxins (e.g. IAA and NAA). Auxin-like activity of kresoxim-Me was also found when stem explants of tobacco were cultured on a hormone-free medium. Kresoxim-Me stimulated shoot formation, particularly at 10<sup>-7</sup> M. The same effect was induced by 10<sup>-8</sup> M IAA. The determination of phytohormone-like substances in shoots of wheat plants foliar-treated with 7 + 10<sup>-4</sup> M kresoxim-Me revealed only slightly changed levels of endogenous IAA, gibberellins and abscisic acid. In contrast, the contents of dihydrozeatin riboside-type cytokinins increased to 160% of the control, while trans-zeatin riboside- and isopentenyladenosine-type cytokinins remained nearly unchanged. The most remarkable alterations were the redns. in 1-aminocyclopropane-1-carboxylic acid (ACC) levels and ethylene formation which were demonstrated in intact plants, leaf disks and the shoots of wheat subjected to drought stress. Kresoxim-Me affected the induction of ACC synthase activity which converts S-adenosyl-methionine to ACC in ethylene biosynthesis. In shoots from foliar-treated wheat plants, 10<sup>-4</sup> M kresoxim-Me inhibited stress-induced increases in endogenous ACC synthase activity, ACC levels and ethylene formation by approx. 50%. Redns. in ACC synthase activity and ACC levels of 30% were also obtained at low concns. of  $\alpha$ -NAA (10<sup>-6</sup> M). In contrast, ACC synthase activity in vitro was not influenced by adding the compds. In wheat leaf disks, the inhibiting effect of kresoxim-Me,  $\alpha$ -NAA and IAA on ethylene formation was accompanied by delayed leaf senescence, characterized by reduced chlorophyll loss. However, in contrast to kresoxim-Me which showed only inhibitory activity on ethylene synthesis over a wide range of concns. applied, the auxins stimulated ethylene production at high concns. of about 10<sup>-4</sup> M. The inhibition of ethylene biosynthesis by kresoxim-Me, together with an increase in endogenous cytokinins could explain the



retardation of senescence and the intensified green leaf  
pigmentation in wheat exposed to this strobilurin.  
REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE  
FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

L11 ANSWER 7 OF 18 CAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 1997:128560 CAPLUS Full-text  
DOCUMENT NUMBER: 126:140904  
ORIGINAL REFERENCE NO.: 126:27135a,27138a  
TITLE: Inhibition of methionine biosynthesis in  
Botrytis cinerea by the anilino pyrimidine  
fungicide pyrimethanil  
AUTHOR(S): Fritz, Rene; Lanen, Catherine; Colas,  
Virginie;  
Leroux, Pierre  
CORPORATE SOURCE: Institut National de la Recherche Agronomique,  
Unite  
de Phytopharmacie et des Mediateurs Chimiques,  
Versailles, 78026, Fr.  
SOURCE: Pesticide Science (1997), 49(1), 40-46  
CODEN: PSSCBG; ISSN: 0031-613X  
PUBLISHER: Wiley  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB When mycelium of *B. cinerea* was treated with low concns. of  
pyrimethanil, the total amount of free amino acids increased.  
Qual. variations were also induced: alanine, glutamine, lysine,  
glycine, histidine, asparagine, arginine, threonine,  $\alpha$ -  
aminobutyrate and  $\beta$ -alanine were accumulated; cyst(e)ine, valine,  
leucine and citrulline were reduced. When mycelium of *B. cinerea*  
was incubated with Na<sub>2</sub>[<sup>35</sup>S]O<sub>4</sub>, pyrimethanil, at 1.5  $\mu$ M, induced a  
decrease of [<sup>35</sup>S]methionine and simultaneously an increase of  
[<sup>35</sup>S]cystathionine. Thus, pyrimethanil inhibits the biosynthesis  
of methionine and suggest that the primary target could be the  
cystathionine  $\beta$ -lyase.

REFERENCE COUNT: 23 THERE ARE 23 CITED REFERENCES AVAILABLE  
FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

L11 ANSWER 8 OF 18 CAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 1993:76896 CAPLUS Full-text  
DOCUMENT NUMBER: 118:76896  
ORIGINAL REFERENCE NO.: 118:13411a,13414a  
TITLE: Control of growth and development of  
Ceratocystis  
fimbriata Ell. et Halst. by plant growth  
regulators.  
IV. Ethylene  
AUTHOR(S): Stopinska, Jadwiga; Kuik, Krystyna  
CORPORATE SOURCE: Inst. Biol., N. Copernicus Univ., Torun, 87-  
100, Pol.  
SOURCE: Bulletin of the Polish Academy of Sciences:  
Biological Sciences (1991), 39(3), 291-300

CODEN: BPABEN; ISSN: 0239-751X

DOCUMENT TYPE:

Journal

LANGUAGE:

English

AB C. fibriata was cultured on potato-dextrose agar on liquid medium containing 2-chloroethylphosphonic acid (CEPA), an ethylene-releasing compound, at 10<sup>-6</sup>-10<sup>-3</sup> M concns. Ethylene inhibited growth of the fungus, sporulation and spore germination. The inhibition was stronger at higher concns. of ethylene. The older mycelium was more sensitive to ethylene than the younger one. C. fibriata produced ethylene enzymically in the presence and also without methionine in the medium. The younger (nonsporulating) mycelium with the high growth intensity produced more ethylene than the sporulating and older mycelium. The fungus did not produce ethylene nonenzymically after 24 h from killing of mycelium.

L11 ANSWER 9 OF 18 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1991:160762 CAPLUS Full-text

DOCUMENT NUMBER: 114:160762

ORIGINAL REFERENCE NO.: 114:27103a,27106a

TITLE:  $\beta$ -(3-Isoxazolin-5-on-2-yl)-alanine from Pisum: allelopathic properties and antimycotic

bioassay

AUTHOR(S):

Schenk, Sigrid U.; Werner, Dietrich

CORPORATE SOURCE:

Bot. Inst., Philipps-Univ. Marburg, Marburg,

D-3550,

Germany

SOURCE:

Phytochemistry (1991), 30(2), 467-70

CODEN: PYTCAS; ISSN: 0031-9422

DOCUMENT TYPE:

Journal

LANGUAGE:

English

AB Grasses and Lactuca sativa when germinated in the presence of the non-protein amino acid  $\beta$ -(3-isoxazolin-5-on-2-yl)-alanine ( $\beta$ IA) from roots and root exudates of pea (P. sativum) seedlings, showed a pronounced reduction of root length and a necrosis of the root tips. Growth of legume seedlings was only slightly affected, indicating the role of this secondary plant product as an allelochem. Besides its effect on plant morphogenesis,  $\beta$ IA also exhibits an antimycotic activity towards Saccharomyces cerevisiae with a min. inhibitory concentration (MIC) of 0.5 ppm.

L11 ANSWER 10 OF 18 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1986:181661 CAPLUS Full-text

DOCUMENT NUMBER: 104:181661

ORIGINAL REFERENCE NO.: 104:28673a,28676a

TITLE: Protection of wheat seedlings from Helminthosporium

infection by seed treatment with chemicals

AUTHOR(S):

Hait, G. N.; Sinha, A. K.

CORPORATE SOURCE:

Dep. Plant Pathol., Bidhan Chandra Krishi

Viswavidyalaya, Kalyani, 741235, India

SOURCE:

Journal of Phytopathology (1986), 115(2), 97-

107

CODEN: JPHYEB; ISSN: 0931-1785

DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Of 24 phytoalexin-inducing chems. studied, HgCl<sub>2</sub>, CuCl<sub>2</sub>, and CdCl<sub>2</sub> totally inhibited the germination of *H. sativum*; Ni(NO<sub>3</sub>)<sub>2</sub>, Na selenite, cycloheximide, IAA [87-51-4] and 2,4-D [94-75-7] inhibited spore germination by 79, 66, 68, 52, and 54%, resp. A few compds. such as DL-norvaline [760-78-1] and DL-methionine [59-51-8] stimulated spore germination. Most compds. when applied in seed treatments effectively protected 3-wk-old wheat seedlings against *H. sativum* infection.

L11 ANSWER 11 OF 18 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1985:144713 CAPLUS Full-text

DOCUMENT NUMBER: 102:144713

ORIGINAL REFERENCE NO.: 102:22647a,22650a

TITLE: Studies on the mode of action of cymoxanil

AUTHOR(S): Fritz, R.; Despreaux, D.; Leroux, P.

CORPORATE SOURCE: Lab. Phytopharm., Inst. Natl. Rech. Agron., Versailles, F-78000, Fr.

SOURCE: Tagungsbericht - Akademie der Landwirtschaftswissenschaften der Deutschen Demokratischen Republik (1984), 222(Syst.

Fungic.

Antifungal Compd.), 65-9

CODEN: TALDA3; ISSN: 0138-2659

DOCUMENT TYPE: Journal

LANGUAGE: English

AB In *Botrytis cinerea*, cymoxanil (I) [57966-95-7] inhibited mycelial growth, and to a lesser extent spore germination. The toxicity of I to *B. cinerea* was antagonized by methionine [63-68-3], glycine [56-40-6], serine [56-45-1], and cysteine [52-90-4]. I transiently inhibited the respiration of *B. cinerea* and *Phytophthora cinnamomi*. I enhanced the incorporation of acetate-14C into lipids in *B. cinerea*, but had a reverse effect in *P. cinnamomi*. I inhibited the penetration and incorporation of uridine-14C, serine-14C, and L-phenylalanine-14C, in both species.

L11 ANSWER 12 OF 18 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1984:116335 CAPLUS Full-text

DOCUMENT NUMBER: 100:116335

ORIGINAL REFERENCE NO.: 100:17613a,17616a

TITLE: Mode of action of hymexazol in *Pythium*

*aphanidermatum*

AUTHOR(S): Nakanishi, Toshiro; Sisler, Hugh D.

CORPORATE SOURCE: Agric. Chem. Res. Lab., Sankyo Co., Shiga, 520-23,

Japan

SOURCE: Nippon Noyaku Gakkaishi (1983), 8(2), 173-81

CODEN: NNGADV; ISSN: 0385-1559

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The dry weight increase of *P. aphanidermatum* hyphae in liquid culture was not affected by exposure to 3 µg/mL hymexazol (I) [10004-44-1] for 5 h, but was almost completely inhibited after this period. Expansion of growing *P. aphanidermatum* hyphae was

inhibited after exposure to 3 µg/mL I for 3 h. Incorporation of 14C of AcONa-2-14C and methionine-methyl-14C into lipids and that of 14C of phenylalanine-U-14C into proteins were not inhibited by 3 µg/mL I during the 1st 3 h of exposure of fungal hyphae. Incorporation of 3H of uridine-6-3H into RNA was inhibited by .apprx.50% after hyphal exposure to I for 3 h, but incorporation of 14C of labeled aspartic acid into RNA and proteins was not inhibited during 3 h of hyphal exposure. I did not interfere with nuclear division or nuclear movement in germinating zoospores. Amts. of I taken up by the fungus reached a maximum within 2 h, indicating that delayed toxicity was not attributable to the time required for I uptake. I was possibly converted into an active derivative inhibitory to a major metabolic pathway, or I directly inhibited an obscure pathway affecting growth only after appreciable delay.

L11 ANSWER 13 OF 18 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1983:48529 CAPLUS Full-text

DOCUMENT NUMBER: 98:48529

ORIGINAL REFERENCE NO.: 98:7423a,7426a

TITLE: Ethylene formation in barley seedlings during early

stages of infection by Drechslera graminea and

its

regulation by seed treatment

AUTHOR(S): Walther, H. F.; Hoffmann, G. M.

CORPORATE SOURCE: Tech. Univ. Muenchen, Freising, D-8050, Fed.

Rep. Ger.

SOURCE: Zeitschrift fuer Pflanzenkrankheiten und

Pflanzenschutz (1982), 89(10), 561-70

CODEN: ZPFPAA; ISSN: 0340-8159

DOCUMENT TYPE: Journal

LANGUAGE: German

AB Natural infection of barley seed with *D. graminea* increased the ethylene [74-85-1] evolution within 3 wk of germination at 2 or 4° from 3-4 to 22 and 12 pmol/mL head space, resp. Seed dressing with Panocrine UTB (I) [74725-91-0] halved the ethylene evolution by infected seedlings. In another test at 4°, dressing with I, ROP 17,660 B (iprodione-carbendazim) [58784-20-6], BAS 39503 F [80123-72-4], or Baytan U [74725-94-3] decreased the ethylene evolution to uninfected control level. Arbosan UT [73730-31-1] And Drawigran plus [84069-57-8] inhibited the ethylene evolution more effectively than did triforine [26644-46-2]. In contrast, Ceresan [107-27-7] increased the ethylene evolution to 110 pmol/mL, evidently by a Hg-induced stress. At 2° all fungicides, with exception of Ceresan and triforine, decreased the ethylene evolution to control levels. Only Ceresan stimulated ethylene formation by noninfected barley. An addition of 10-3 mol L-methionine [63-68-3]/L medium within 4 h induced ethylene evolution by *D. graminea* in vitro, whereas the precursor, α-ketoglutaric acid [328-50-7], was almost ineffective. The usefulness of ethylene evolution tests for fungicide screening is discussed.

L11 ANSWER 14 OF 18 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 1980:632482 CAPLUS Full-text  
 DOCUMENT NUMBER: 93:232482  
 ORIGINAL REFERENCE NO.: 93:37099a,37102a  
 TITLE: Effect of chemical agents on the  
 interrelations between potato plants and Phytophthora  
 infestans (Mont.) D By. III. Effect of  
 organophosphorus pesticides  
 AUTHOR(S): Mustafa, M.; D'yakov, Yu. T.  
 CORPORATE SOURCE: Mosk. Gos. Univ., Moscow, USSR  
 SOURCE: Mikologiya i Fitopatologiya (1980), 14(1), 31-  
 6  
 CODEN: MIFIB2; ISSN: 0026-3648  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Russian  
 GI



AB Preplant treatment of potato tubers with 5-100 µg Cidial [2597-03-7]/mL induced formation of 50-60 µg rishitin [18178-54-6]/mL tuber on contact with *P. infestans* zoospores. Phosalone [2310-17-0], phthalophos [732-11-6], and Sayfos [78-57-9] were less effective. I; R = H, R1 = SP(:S)(OEt)2 [57779-12-1], I; R = H, R1 = P(:O)(OEt)2 [61704-85-6], I; R = Me, R1 = P(:O)(OEt)2 [74748-28-0], and I; R = Me, R1 = P(:O)(OPr)2 [74754-52-2] also induced rishitin formation by the infected tubers and were highly toxic for *P. infestans* zoospores in vitro, whereas O,O-diethyldithiophosphoric acid [298-06-6] failed to stimulate the rishitin formation in spite of its high toxicity for the zoospores in vitro. Quinosan [82-68-8], Inezin [21722-85-0], and ketazin [13286-32-3] induced rishitin formation in infected (but not in healthy) tubers, whereas Pyrazophos [13457-18-6] inhibited rishitin formation in infected tubers, while showing a high toxicity for zoospores in vitro. Inezin, ketazin P [26087-47-8], and Quinosan rapidly stimulated protein and amino acid release from germinating zoospores. Ca(NO3)2 at 50 µg/mL protected the germinating zoospores from protein loss caused by Quinosan. Methionine [63-68-3] and cysteine [52-90-4] were less effective protectants. Ca2+ protected the germinating zoospores from the release of substances which induce rishitin formation in the presence of Quinosan and Inezin.

L11 ANSWER 15 OF 18 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 1979:450986 CAPLUS Full-text

DOCUMENT NUMBER: 91:50986  
ORIGINAL REFERENCE NO.: 91:8215a,8218a  
TITLE: Studies on the inhibitory effects of N-acylamino acid  
bacteria and its analog for the pathogenic fungus and  
in various plants  
AUTHOR(S): Takano, Saburo  
CORPORATE SOURCE: Dep. Agric. Chem., Tokyo Univ. Agric., Tokyo,  
Japan  
SOURCE: Memoirs of the Tokyo University of Agriculture  
(1978),

20, 51-73  
CODEN: TOAMB6; ISSN: 0372-0322  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB N-acyl amino acids were synthesized and their inhibitory effects on pathogenic fungi studied. N-Benzoyl-L-leucine (I) [1466-83-7] and N-phenylacetyl-L-leucine [730-15-4] at 10 mM inhibited the growth of *Rhizoctonia solani* and N-benzoyl-L-methionine [10290-61-6] and N-phenoxyacetyl-L-leucine [14231-46-0] inhibited proliferation of *Pyricularia oryzae*. I inhibited the proliferation of *Gloeosporium musarum* and *Alternaria kikuchiana*. Na-cocoyl-L-arginine Et ester-D,L-2-pyrrolidone 5-carboxylic acid salt (II) at 10 µg/mL controlled (96.4%) *Uromyces fabae* and had a broader and more significant inhibitory effect on spore germination. I or II (100 µg/mL) inhibited *G. musarum* on banana. II inhibited the growth of *Botrytis fabae*, *Gymnosporangium haraeae*, *Venturia nashicola*, and *A. kikuchiana* in pears. II 500-1000, Cu hydroxide chloride 1470, and 8-hydroxyquinolinatocopper [10380-28-6] 772 µg/mL inhibited *Pseudoperonospora cubensis*, *Sphaerotheca fuliginea*, and *Pseudomonas lachrymans* in cucumber. The inhibitory mechanism of II on the growth of pathogenic bacilli includes leakage of biotin, glucose, ATP, and protein from the bacilli.

L11 ANSWER 16 OF 18 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1973:512267 CAPLUS Full-text  
DOCUMENT NUMBER: 79:112267  
ORIGINAL REFERENCE NO.: 79:18206h,18207a  
TITLE: Effect of antimetabolites and fungicides on  
elongation of germination hyphae of powdery  
mildew in vitro  
AUTHOR(S): Van't Land, B. G.; Dekker, J.  
CORPORATE SOURCE: Lab. Phytopathol., Agric. Univ., Wageningen,  
Neth.  
SOURCE: Netherlands Journal of Plant Pathology (1972),  
78(6),

242-6  
CODEN: NJPPAM; ISSN: 0028-2944  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB In vitro germ tube elongation of *Sphaerotheca fuliginea* was inhibited by low fungicide concns. and by high concns. of L-methionine [63-68-3] and procaine-hydrochloride (I) [51-05-8]. D-methionine [348-67-4] was inactive, both in vivo and in vitro. 6-

Azaauracil [461-89-2] was converted to 6-azauridine monophosphate [2018-19-1] by *S. fuliginea*. The effects on hyphal elongation were used in the screening of fungicides and antimetabolites for control of powdery mildew.

L11 ANSWER 17 OF 18 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1964:496311 CAPLUS [Full-text](#)  
DOCUMENT NUMBER: 61:96311  
ORIGINAL REFERENCE NO.: 61:13822g-h  
TITLE: Modes of action of chemotherapeutic agents in plants.  
Discussion  
AUTHOR(S): Cowling, Ellis B.; et al.  
CORPORATE SOURCE: Conn. Agr. Expt. Sta., New Haven  
SOURCE: Conn. Agr. Expt. Sta., New Haven, Bull. No. (1963),

663, 72-7

DOCUMENT TYPE: Journal  
LANGUAGE: Unavailable

AB Chemical differences between pathogens and their plant hosts are considered, with some apparently new data. Relations between phenols and carbohydrate metabolism are discussed. In expts. on fusiform rust (a major disease of southern pine trees), the stem-invading fungus produces stem galls. Cycloheximide (I) in very low concns. prevented the germination of rust spores. I was translocated in slash pine seedlings at concns. high enough to inhibit a test-assay organism (not named) but had no apparent effect on the fungus in the tissue of the infected host. It is possible that I did not diffuse to the sites of infection rapidly enough to affect the pathogen. The relative fungicidal concns. of ethionine (II) on agar (test fungus not named) were 25, 50, and over 1000 p.p.m. for the L-, DL-, and D-forms, resp. Possibly II acted as a competitive inhibitor for methionine required as a Me donor in the formation of pectin. Applications of HgCl<sub>2</sub> or CuCl<sub>2</sub> to the endocarp of pea pods induced the formation of pisatin in concns. which inhibited some pathogens of peas in vitro. Other chemical compds. induced the formation of lower concns. of pisatin.

L11 ANSWER 18 OF 18 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1960:103999 CAPLUS [Full-text](#)  
DOCUMENT NUMBER: 54:103999  
ORIGINAL REFERENCE NO.: 54:19837h-i,19838a-b  
TITLE: Reversal of fungitoxicity of 8-quinolinol by amino

acids and other chelators

AUTHOR(S): Zentmyer, George A.; Rich, Saul; Horsfall, James G.  
CORPORATE SOURCE: Univ. of California, Riverside  
SOURCE: Phytopathology (1960), 50, 421-4  
CODEN: PHYTAJ; ISSN: 0031-949X

DOCUMENT TYPE: Journal  
LANGUAGE: Unavailable

AB Amino acids and other chelators were tested for their ability to reverse the toxicity of 8-quinolinol (I) to spores and mycelium of

Aspergillus niger and to mycelium of Botryosphaeria ribis. In the spore test with A. niger, cysteine, histidine, tryptophan, Casamino acids, dithizone, and versene reversed I toxicity, and glutamic acid, asparagine, and glutathione did not. In the mycelium test with A. niger, cysteine reversed I toxicity; glutathione, asparagine, histidine, and tryptophan had no effect, and glutamic acid increased the toxicity of I. In the test with B. ribis, cysteine reversed I toxicity, and glutathione, asparagine, histidine, tryptophan, glutamic acid, glycine, and methionine had no effect. Dithizone and quinaldic acid reversed the toxicity of I to the spores of Stemphylium sarciniforme and Monilinia fructicola. In vitro studies showed that 0.5% solns. of histidine and cysteine can remove Cu from a 7 p.p.m. solution of Cu oxinate (II). It is suggested that II produces fungitoxicity in the following manner. The amino acids of the cell take Cu from the half-chelated II, and release I in situ. The Cu poisons amino acids, proteins, and enzymes while the freed I sequesters prosthetic trace metals such as Fe++, Zn++, and Co++.

```
=> s (fungicid? or anti!fung? or pesticid?) and (methionine) and
(mycelium or germinat?)
  118517 FUNGICID?
    3 ANTI!FUNG?
  98639 PESTICID?
  97295 METHIONINE
    557 METHIONINES
  97489 METHIONINE
    (METHIONINE OR METHIONINES)
  16266 MYCELIUM
    29 MYCELIUMS
    9113 MYCELIA
      2 MYCELIAS
  23446 MYCELIUM
    (MYCELIUM OR MYCELIUMS OR MYCELIA OR MYCELIAS)
  63685 GERMINAT?
L12 29 (FUNGICID? OR ANTI!FUNG? OR PESTICID?) AND (METHIONINE)
AND (MYC
      ELIUM OR GERMINAT?)
```

```
=> s l12 and (py,2003 or ay<2003 or pry<2003)
  17516 PY
    771 PIES
  18286 PY
    (PY OR PIES)
  42924 2003
    0 PY,2003
      (PY(W)2003)
  4503738 AY<2003
  3972615 PRY<2003
L13 1 L12 AND (PY,2003 OR AY<2003 OR PRY<2003)
```

```
=> d l13 ibib abs
```

```
L13 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2009 ACS on STN
```



ACCESSION NUMBER: 2004:269834 CAPLUS Full-text  
 DOCUMENT NUMBER: 140:266136  
 TITLE: Seed treatment for germination stimulation  
 and plant vigor enhancement  
 INVENTOR(S): Johnson, William S.  
 PATENT ASSIGNEE(S): USA  
 SOURCE: U.S. Pat. Appl. Publ., 10 pp.  
 CODEN: USXXCO  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20040063582	A1	20040401	US 2002-246093	
20020917 <--				
US 7001869	B2	20060221		
PRIORITY APPLN. INFO.:			US 2002-246093	
20020917 <--				
AB A seed treatment composition is given, containing plant macronutrients, micronutrients, a pesticides and at least one growth regulator. The composition addnl. contains vitamins, amino acids, penetrants and an energy source. The treatment results in germination stimulation and plant vigor and hardness enhancement.				
REFERENCE COUNT:	14	THERE ARE 14 CITED REFERENCES AVAILABLE		
FOR THIS				
RECORD. ALL CITATIONS AVAILABLE IN THE				
RE FORMAT				

```

=> s (fungicid? or anti!fung? or pesticid? or herbici?) and
(methionine) and (mycelium or germinat?)
  118517 FUNGICID?
    3 ANTI!FUNG?
  98639 PESTICID?
  93945 HERBICI?
  97295 METHIONINE
    557 METHIONINES
  97489 METHIONINE
    (METHIONINE OR METHIONINES)
  16266 MYCELIUM
    29 MYCELIUMS
    9113 MYCELIA
      2 MYCELIAS
  23446 MYCELIUM
    (MYCELIUM OR MYCELIUMS OR MYCELIA OR MYCELIAS)
  63685 GERMINAT?
L14 36 (FUNGICID? OR ANTI!FUNG? OR PESTICID? OR HERBICI?) AND
(METHIONI
      NE) AND (MYCELIUM OR GERMINAT?)

=> s l14 and (py<2003 or ay<2003 or pry<2003)
  22983274 PY<2003
  4503738 AY<2003
  3972615 PRY<2003
  
```

L15

27 L14 AND (PY<2003 OR AY<2003 OR PRY<2003)

=> d l15 ibib abs 1-10

L15 ANSWER 1 OF 27 CAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 2004:269834 CAPLUS Full-text  
DOCUMENT NUMBER: 140:266136  
TITLE: Seed treatment for germination stimulation  
and plant vigor enhancement  
INVENTOR(S): Johnson, William S.  
PATENT ASSIGNEE(S): USA  
SOURCE: U.S. Pat. Appl. Publ., 10 pp.  
CODEN: USXXCO  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

	PATENT NO. -----	KIND ----	DATE -----	APPLICATION NO. -----	DATE ----
----	US 20040063582	A1	20040401	US 2002-246093	
20020917 <--	US 7001869	B2	20060221		
PRIORITY APPLN. INFO.:				US 2002-246093	
20020917 <--					
AB	A seed treatment composition is given, containing plant macronutrients, micronutrients, a pesticides and at least one growth regulator. The composition addnl. contains vitamins, amino acids, penetrants and an energy source. The treatment results in germination stimulation and plant vigor and hardness enhancement.				
REFERENCE COUNT:	14	THERE ARE 14 CITED REFERENCES AVAILABLE			
FOR THIS					

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

L15 ANSWER 2 OF 27 CAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 2001:627332 CAPLUS Full-text  
DOCUMENT NUMBER: 135:299904  
TITLE: Nitrolin and techniques for its use on winter  
wheat  
crops  
AUTHOR(S): Niyazmetov, U. K.; Kariev, A. U.;  
Dustmukhamedov, T.  
CORPORATE SOURCE: Inst. Khim. Rastitel'nykh Veshchestv im. S.  
Yu.  
Yunusova, AN Ruz, Uzbekistan  
SOURCE: Doklady Akademii Nauk Respubliki Uzbekistan ( 2001), (3), 34-37  
CODEN: DARUEE; ISSN: 1019-8954  
PUBLISHER: Fan  
DOCUMENT TYPE: Journal  
LANGUAGE: Russian  
AB Growth regulating activity of nitrolin and its compatibility with  
seed treatment with the fungicide tuzal were investigated. The  
following parameters were used: seed germination, susceptibility

to root rot, grain yield, content of starch in the grain, and content of selected amino acids in the leaves. As a result of growth processes intensification, the plants treated with nitrolin had higher rate of germination compared to control plants, lower number of diseased plants, higher grain yield and higher starch content in the grain. The decrease in root rot occurrence was also supplemented by the fungicidal action of tuzal. The composition of amino acids in treated plants did not differ from the control, although their content was higher in leaves of nitrolin treated plants.

L15 ANSWER 3 OF 27 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1998:121042 CAPLUS Full-text

DOCUMENT NUMBER: 128:163907

ORIGINAL REFERENCE NO.: 128:32199a,32202a

TITLE: The effect of herbicides applied at different terms on protein content and amino acids composition in the winter wheat grain of the

Arda and

Juma varieties

AUTHOR(S): Ostapczuk, Elzbieta; Rola, Henryka; Sykut, Anna;

Nowicka, Barbara

CORPORATE SOURCE: Akademia Rolnicza, Lublin, 20-950, Pol.

SOURCE: Pestycydy (Warsaw) (1997), (1-2), 59-65

CODEN: PSTYDL; ISSN: 0208-8703

PUBLISHER: Instytut Przemyslu Organicznego

DOCUMENT TYPE: Journal

LANGUAGE: Polish

AB The 3 yr field experiment (1993-1995) studied the effect of Dicuran 80 WP, Glean 75 DF, Quartz Super, Grodyl 75 WG, Racer 25 EC and Stomp 330 EC on the content of protein and 16 amino acids in the grain of winter wheat of the varieties Arda and Juma. Before germination (I term), after germination in autumn (II term) and in spring (III term) herbicides were applied. The effect of the herbicides was only slight and it was related to the wheat variety. Dicuran and Glean decreased total protein and aspartic acid content; Grodyl increased protein content in the variety Arda. Glean, Racer, Stomp decreased, and Dicuran, Quartz Super increased protein content in the variety Juma. In this variety, Stomp and Quartz Super increased aspartic acid, glutamic acid, leucine, methionine and threonine content.

L15 ANSWER 4 OF 27 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1997:380005 CAPLUS Full-text

DOCUMENT NUMBER: 127:1954

ORIGINAL REFERENCE NO.: 127:463a,466a

TITLE: Bioregulatory effects of the fungicidal strobilurin kresoxim-methyl in wheat (Triticum aestivum)

AUTHOR(S): Grossmann, Klaus; Retzlaff, Gunter

CORPORATE SOURCE: Agricultural Res. Station, BASF, Limbergerhof, D-67114, Germany

SOURCE: Pesticide Science (1997), 50(1), 11-20

CODEN: PSSCBG; ISSN: 0031-613X

PUBLISHER: Wiley

DOCUMENT TYPE: Journal

LANGUAGE: English

AB    Apart from its fungicidal effect, the strobilurin kresoxim-Me (BAS 490 F) was found to induce physiol. and developmental alterations in wheat (*Triticum aestivum* L.) which are seen in connection with improved yield. In a series of biotests including heterotrophic maize and photoautotrophic algal cell suspensions, duckweed, isolated mustard shoots and germinating cress seeds, kresoxim-Me showed a similar response pattern to standard auxins (e.g. IAA and NAA). Auxin-like activity of kresoxim-Me was also found when stem explants of tobacco were cultured on a hormone-free medium. Kresoxim-Me stimulated shoot formation, particularly at 10<sup>-7</sup> M. The same effect was induced by 10<sup>-8</sup> M IAA. The determination of phytohormone-like substances in shoots of wheat plants foliar-treated with 7 + 10<sup>-4</sup> M kresoxim-Me revealed only slightly changed levels of endogenous IAA, gibberellins and abscisic acid. In contrast, the contents of dihydrozeatin riboside-type cytokinins increased to 160% of the control, while trans-zeatin riboside- and isopentenyladenosine-type cytokinins remained nearly unchanged. The most remarkable alterations were the redns. in 1-aminocyclopropane-1-carboxylic acid (ACC) levels and ethylene formation which were demonstrated in intact plants, leaf disks and the shoots of wheat subjected to drought stress. Kresoxim-Me affected the induction of ACC synthase activity which converts S-adenosyl-methionine to ACC in ethylene biosynthesis. In shoots from foliar-treated wheat plants, 10<sup>-4</sup> M kresoxim-Me inhibited stress-induced increases in endogenous ACC synthase activity, ACC levels and ethylene formation by approx. 50%. Redns. in ACC synthase activity and ACC levels of 30% were also obtained at low concns. of  $\alpha$ -NAA (10<sup>-6</sup> M). In contrast, ACC synthase activity in vitro was not influenced by adding the compds. In wheat leaf disks, the inhibiting effect of kresoxim-Me,  $\alpha$ -NAA and IAA on ethylene formation was accompanied by delayed leaf senescence, characterized by reduced chlorophyll loss. However, in contrast to kresoxim-Me which showed only inhibitory activity on ethylene synthesis over a wide range of concns. applied, the auxins stimulated ethylene production at high concns. of about 10<sup>-4</sup> M. The inhibition of ethylene biosynthesis by kresoxim-Me, together with an increase in endogenous cytokinins could explain the retardation of senescence and the intensified green leaf pigmentation in wheat exposed to this strobilurin.

REFERENCE COUNT:        28        THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

L15 ANSWER 5 OF 27 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER:        1997:128560 CAPLUS Full-text

DOCUMENT NUMBER:        126:140904

ORIGINAL REFERENCE NO.: 126:27135a,27138a

TITLE:                    Inhibition of methionine biosynthesis in Botrytis cinerea by the anilinopyrimidine fungicide pyrimethanil

AUTHOR(S):                Fritz, Rene; Lanen, Catherine; Colas,

Virginie;

CORPORATE SOURCE:  
Unite

Leroux, Pierre  
Institut National de la Recherche Agronomique,

de Phytopharmacie et des Mediateurs Chimiques,  
Versailles, 78026, Fr.

SOURCE: Pesticide Science (1997), 49(1), 40-46  
CODEN: PSSCBG; ISSN: 0031-613X

PUBLISHER: Wiley  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB When mycelium of *B. cinerea* was treated with low concns. of pyrimethanil, the total amount of free amino acids increased. Qual. variations were also induced: alanine, glutamine, lysine, glycine, histidine, asparagine, arginine, threonine,  $\alpha$ -aminobutyrate and  $\beta$ -alanine were accumulated; cyst(e)ine, valine, leucine and citrulline were reduced. When mycelium of *B. cinerea* was incubated with Na<sub>2</sub>[<sup>35</sup>S]O<sub>4</sub>, pyrimethanil, at 1.5  $\mu$ M, induced a decrease of [<sup>35</sup>S]methionine and simultaneously an increase of [<sup>35</sup>S]cystathionine. Thus, pyrimethanil inhibits the biosynthesis of methionine and suggest that the primary target could be the cystathionine  $\beta$ -lyase.

REFERENCE COUNT: 23 THERE ARE 23 CITED REFERENCES AVAILABLE  
FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

L15 ANSWER 6 OF 27 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1993:76896 CAPLUS Full-text

DOCUMENT NUMBER: 118:76896

ORIGINAL REFERENCE NO.: 118:13411a,13414a

TITLE: Control of growth and development of

*Ceratocystis*

fimbriata Ell. et Halst. by plant growth

regulators.

IV. Ethylene

AUTHOR(S): Stopinska, Jadwiga; Kuik, Krystyna

CORPORATE SOURCE: Inst. Biol., N. Copernicus Univ., Torun, 87-100, Pol.

SOURCE: Bulletin of the Polish Academy of Sciences:  
Biological Sciences (1991), 39(3), 291-300  
CODEN: BPABEN; ISSN: 0239-751X

DOCUMENT TYPE: Journal  
LANGUAGE: English

AB *C. fimbriata* was cultured on potato-dextrose agar on liquid medium containing 2-chloroethylphosphonic acid (CEPA), an ethylene-releasing compound, at 10<sup>-6</sup>-10<sup>-3</sup> M concns. Ethylene inhibited growth of the fungus, sporulation and spore germination. The inhibition was stronger at higher concns. of ethylene. The older mycelium was more sensitive to ethylene than the younger one. *C. fimbriata* produced ethylene enzymically in the presence and also without methionine in the medium. The younger (nonsporulating) mycelium with the high growth intensity produced more ethylene than the sporulating and older mycelium. The fungus did not produce ethylene nonenzymically after 24 h from killing of mycelium.

L15 ANSWER 7 OF 27 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 1992:124848 CAPLUS Full-text  
 DOCUMENT NUMBER: 116:124848  
 ORIGINAL REFERENCE NO.: 116:21013a,21016a  
 TITLE: Broad antifungal activity of  
 $\beta$ -isoxazolinonyl-alanine, a non-protein amino  
 acid from roots of pea (*Pisum sativum* L.)  
 seedlings  
 AUTHOR(S): Schenk, S. U.; Lambein, F.; Werner, D.  
 CORPORATE SOURCE: Bot. Inst., Philipps-Univ., Marburg, W-3550,  
 Germany  
 SOURCE: Biology and Fertility of Soils (1991),  
 11(3), 203-9  
 CODEN: BFSOEE; ISSN: 0178-2762  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB  $\beta$ -(Isioxazolin-5-on-2yl)alanine ( $\beta$ IA), a heterocyclic nonprotein  
 amino acid from root exts. and root exudates of pea seedlings,  
 acts as a potent growth inhibitor of several eukaryotic organisms,  
 including yeasts, phytopathogenic fungi, unicellular green algae,  
 and higher plants. The antibiotic effect on bakers' yeast was  
 reversed by L-methionine, L-cysteine, and L-homocysteine.  
 Phytopathogenic fungi such as *Botrytis cinerea*, *Pythium ultimum*,  
 and *Rhizoctonia solani* grown on agar containing  $\beta$ IA were inhibited  
 in the growth of mycelia or in the production of sclerotia. In  
 contrast, no significant inhibition of either gram-pos. or gram-  
 neg. bacteria was observed *Rhizobium leguminosarum*, the  
 compatible microsymbiont of *Pisum* spp., and *Rhizobium meliloti*  
 tolerated  $\leq 2.9$  mM  $\beta$ IA (500 ppm) without affecting the growth rate.  
*Bradyrhizobium japonicum* even gave a pos. chemotactic response to  
 $\beta$ IA. The ecol. significance of  $\beta$ IA as a preformed plant  
 protectant during the seedling stage of *Pisum* spp. and other  $\beta$ IA-  
 containing legumes is discussed.

L15 ANSWER 8 OF 27 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 1991:673298 CAPLUS Full-text  
 DOCUMENT NUMBER: 115:273298  
 ORIGINAL REFERENCE NO.: 115:46285a,46288a  
 TITLE: Application of NMR spectrometry to fungicide  
 pharmacology  
 AUTHOR(S): Yoshida, Mitsuru  
 CORPORATE SOURCE: Natl. Inst. Agro-Environ. Sci., Tsukuba, 305,  
 Japan  
 SOURCE: Nippon Noyaku Gakkaishi (1991), 16(3),  
 545-54  
 CODEN: NNGADV; ISSN: 0385-1559  
 DOCUMENT TYPE: Journal; General Review  
 LANGUAGE: Japanese  
 AB A review with 51 refs., on the author's work on the title subject,  
 in which  $^{13}\text{C}$  and  $^1\text{H}$  NMR were applied to elucidate fungicidal  
 action on transmethylation from methionine to choline in fungal  
 mycelia and on water permeability of fungal cell membrane, resp.,

and two-dimensional 1H NMR was applied to the anal. of the binding of berenil with DNA.

L15 ANSWER 9 OF 27 CAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 1991:160762 CAPLUS Full-text  
DOCUMENT NUMBER: 114:160762  
ORIGINAL REFERENCE NO.: 114:27103a,27106a  
TITLE:  $\beta$ -(3-Isioxazolin-5-on-2-yl)-alanine from Pisum:  
allelopathic properties and antimycotic  
bioassay  
AUTHOR(S): Schenk, Sigrid U.; Werner, Dietrich  
CORPORATE SOURCE: Bot. Inst., Philipps-Univ. Marburg, Marburg,  
D-3550,  
Germany  
SOURCE: Phytochemistry (1991), 30(2), 467-70  
CODEN: PYTCAS; ISSN: 0031-9422  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB Grasses and Lactuca sativa when germinated in the presence of the non-protein amino acid  $\beta$ -(3-isoxazolin-5-on-2-yl)-alanine ( $\beta$ IA) from roots and root exudates of pea (P. sativum) seedlings, showed a pronounced reduction of root length and a necrosis of the root tips. Growth of legume seedlings was only slightly affected, indicating the role of this secondary plant product as an allelochem. Besides its effect on plant morphogenesis,  $\beta$ IA also exhibits an antimycotic activity towards Saccharomyces cerevisiae with a min. inhibitory concentration (MIC) of 0.5 ppm.

L15 ANSWER 10 OF 27 CAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 1990:34661 CAPLUS Full-text  
DOCUMENT NUMBER: 112:34661  
ORIGINAL REFERENCE NO.: 112:5989a,5992a  
TITLE: Amino acids alterations in stored seeds under stress  
of methyl parathion and lindane dressing. II.  
Wheat  
grains  
AUTHOR(S): Afifi, F. A.; El-Ballal, A. S.  
CORPORATE SOURCE: Fac. Agric., Ain Shams Univ., Cairo, Egypt  
SOURCE: Egyptian Journal of Physiological Sciences (1989), Volume Date 1986, 13(1-2), 123-33  
CODEN: EJPLAD; ISSN: 0301-8660  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB The effect of methyl parathion (0.5 ppm) and lindane (0.1 ppm) on germination and amino acids of stored wheat grains was studied. The pesticides affected the free and more significantly the conjugated amino acids. The effect of the 2 pesticides on different amino acids depended on the type of amino acids.

=> s (spore? and germinat?) and (anti!fung? or fungicid? or pesticid?)

or herbicid?)  
38613 SPORE?  
63685 GERMINAT?  
3 ANTI!FUNG?  
118517 FUNGICID?  
98639 PESTICID?  
93940 HERBICID?  
L16 2047 (SPORE? AND GERMINAT?) AND (ANTI!FUNG? OR FUNGICID? OR  
PESTICID?  
OR HERBICID?)

=> s (spore? and germinat?) and (anti!fung? or fungicid? or pesticid?  
or herbicid?) and (methionine)  
38613 SPORE?  
63685 GERMINAT?  
3 ANTI!FUNG?  
118517 FUNGICID?  
98639 PESTICID?  
93940 HERBICID?  
97295 METHIONINE  
557 METHIONINES  
97489 METHIONINE  
(METHIONINE OR METHIONINES)  
L17 8 (SPORE? AND GERMINAT?) AND (ANTI!FUNG? OR FUNGICID? OR  
PESTICID?  
OR HERBICID?) AND (METHIONINE)

=> s l17 and (py<2003 or ay<2003 or pry<2003)  
22983274 PY<2003  
4503738 AY<2003  
3972615 PRY<2003  
L18 6 L17 AND (PY<2003 OR AY<2003 OR PRY<2003)

=> d l18 ibib abs 1-6

L18 ANSWER 1 OF 6 CAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 1993:76896 CAPLUS Full-text  
DOCUMENT NUMBER: 118:76896  
ORIGINAL REFERENCE NO.: 118:13411a,13414a  
TITLE: Control of growth and development of  
Ceratomyces fimbriata Ell. et Halst. by plant growth  
regulators.  
IV. Ethylene  
AUTHOR(S): Stopinska, Jadwiga; Kuik, Krystyna  
CORPORATE SOURCE: Inst. Biol., N. Copernicus Univ., Torun, 87-  
100, Pol.  
SOURCE: Bulletin of the Polish Academy of Sciences:  
Biological Sciences (1991), 39(3), 291-300  
CODEN: BPABEN; ISSN: 0239-751X  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB C. fimbriata was cultured on potato-dextrose agar on liquid medium  
containing 2-chloroethylphosphonic acid (CEPA), an ethylene-  
releasing compound, at 10-6-10-3 M concns. Ethylene inhibited  
growth of the fungus, sporulation and spore germination. The  
inhibition was stronger at higher concns. of ethylene. The older



mycelium was more sensitive to ethylene than the younger one. C. fibriata produced ethylene enzymically in the presence and also without methionine in the medium. The younger (nonsporulating) mycelium with the high growth intensity produced more ethylene than the sporulating and older mycelium. The fungus did not produce ethylene nonenzymically after 24 h from killing of mycelium.

L18 ANSWER 2 OF 6 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1986:181661 CAPLUS Full-text

DOCUMENT NUMBER: 104:181661

ORIGINAL REFERENCE NO.: 104:28673a,28676a

TITLE: Protection of wheat seedlings from  
Helminthosporium

AUTHOR(S): Hait, G. N.; Sinha, A. K.

CORPORATE SOURCE: Dep. Plant Pathol., Bidhan Chandra Krishi  
Viswavidyalaya, Kalyani, 741235, India

SOURCE: Journal of Phytopathology (1986), 115(2),  
97-107

CODEN: JPHYEB; ISSN: 0931-1785

DOCUMENT TYPE: Journal

LANGUAGE: English

AB OF 24 phytoalexin-inducing chems. studied, HgCl<sub>2</sub>, CuCl<sub>2</sub>, and CdCl<sub>2</sub> totally inhibited the germination of H. sativum; Ni(NO<sub>3</sub>)<sub>2</sub>, Na selenite, cycloheximide, IAA [87-51-4] and 2,4-D [94-75-7] inhibited spore germination by 79, 66, 68, 52, and 54%, resp. A few compds. such as DL-norvaline [760-78-1] and DL-methionine [59-51-8] stimulated spore germination. Most compds. when applied in seed treatments effectively protected 3-wk-old wheat seedlings against H. sativum infection.

L18 ANSWER 3 OF 6 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1985:144713 CAPLUS Full-text

DOCUMENT NUMBER: 102:144713

ORIGINAL REFERENCE NO.: 102:22647a,22650a

TITLE: Studies on the mode of action of cymoxanil

Fritz, R.; Despreaux, D.; Leroux, P.

CORPORATE SOURCE: Lab. Phytopharm., Inst. Natl. Rech. Agron.,  
Versailles, F-78000, Fr.

SOURCE: Tagungsbericht - Akademie der  
Landwirtschaftswissenschaften der Deutschen  
Demokratischen Republik (1984), 222(Syst.  
Fungic. Antifungal Compd.), 65-9

CODEN: TALDA3; ISSN: 0138-2659

DOCUMENT TYPE: Journal

LANGUAGE: English

AB In Botrytis cinerea, cymoxanil (I) [57966-95-7] inhibited mycelial growth, and to a lesser extent spore germination. The toxicity of I to B. cinerea was antagonized by methionine [63-68-3], glycine [56-40-6], serine [56-45-1], and cysteine [52-90-4]. I transiently inhibited the respiration of B. cinerea and Phytophthora cinnamomi. I enhanced the incorporation of acetate-<sup>14</sup>C into lipids in B. cinerea, but had a reverse effect in P.

cinnamomi. I inhibited the penetration and incorporation of uridine-14C, serine-14C, and L-phenylalanine-14C, in both species.

L18 ANSWER 4 OF 6 CAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 1980:632482 CAPLUS Full-text  
DOCUMENT NUMBER: 93:232482  
ORIGINAL REFERENCE NO.: 93:37099a,37102a  
TITLE: Effect of chemical agents on the  
interrelations between potato plants and Phytophthora  
infestans (Mont.) D By. III. Effect of  
organophosphorus pesticides  
AUTHOR(S): Mustafa, M.; D'yakov, Yu. T.  
CORPORATE SOURCE: Mosk. Gos. Univ., Moscow, USSR  
SOURCE: Mikologiya i Fitopatologiya (1980), 14(1),  
31-6  
CODEN: MIFIB2; ISSN: 0026-3648  
DOCUMENT TYPE: Journal  
LANGUAGE: Russian  
GI



AB Preplant treatment of potato tubers with 5-100 µg Cidial [2597-03-7]/mL induced formation of 50-60 µg rishitin [18178-54-6]/mL tuber on contact with *P. infestans* zoospores. Phosalone [2310-17-0], phthalophos [732-11-6], and Sayfos [78-57-9] were less effective. I; R = H, R1 = SP(:S)(OEt)2 [57779-12-1], I; R = H, R1 = P(:O)(OEt)2 [61704-85-6], I; R = Me, R1 = P(:O)(OEt)2 [74748-28-0], and I; R = Me, R1 = P(:O)(OPr)2 [74754-52-2] also induced rishitin formation by the infected tubers and were highly toxic for *P. infestans* zoospores in vitro, whereas O,O-diethyldithiophosphoric acid [298-06-6] failed to stimulate the rishitin formation in spite of its high toxicity for the zoospores in vitro. Quinosan [82-68-8], Inezin [21722-85-0], and ketazin [13286-32-3] induced rishitin formation in infected (but not in healthy) tubers, whereas Pyrazophos [13457-18-6] inhibited rishitin formation in infected tubers, while showing a high toxicity for zoospores in vitro. Inezin, ketazin P [26087-47-8], and Quinosan rapidly stimulated protein and amino acid release from germinating zoospores. Ca(NO3)2 at 50 µg/mL protected the germinating zoospores from protein loss caused by Quinosan. Methionine [63-68-3] and cysteine [52-90-4] were less effective protectants. Ca2+ protected the germinating zoospores from the release of substances which induce rishitin formation in the presence of Quinosan and Inezin.

L18 ANSWER 5 OF 6 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1979:450986 CAPLUS Full-text

DOCUMENT NUMBER: 91:50986

ORIGINAL REFERENCE NO.: 91:8215a,8218a

TITLE: Studies on the inhibitory effects of N-acylamino acid

and its analog for the pathogenic fungus and

bacteria

in various plants

AUTHOR(S): Takano, Saburo

CORPORATE SOURCE: Dep. Agric. Chem., Tokyo Univ. Agric., Tokyo, Japan

SOURCE: Memoirs of the Tokyo University of Agriculture (

1978), 20, 51-73

CODEN: TOAMB6; ISSN: 0372-0322

DOCUMENT TYPE: Journal

LANGUAGE: English

AB N-acyl amino acids were synthesized and their inhibitory effects on pathogenic fungi studied. N-Benzoyl-L-leucine (I) [1466-83-7] and N-phenylacetyl-L-leucine [730-15-4] at 10 mM inhibited the growth of *Rhizoctonia solani* and N-benzoyl-L-methionine [10290-61-6] and N-phenoxyacetyl-L-leucine [14231-46-0] inhibited proliferation of *Pycularia oryzae*. I inhibited the proliferation of *Gloeosporium musarum* and *Alternaria kikuchiana*. Na-cocoyl-L-arginine Et ester-D,L-2-pyrrolidone 5-carboxylic acid salt (II) at 10 µg/mL controlled (96.4%) *Uromyces fabae* and had a broader and more significant inhibitory effect on spore germination. I or II (100 µg/mL) inhibited *G. musarum* on banana. II inhibited the growth of *Botrytis fabae*, *Gymnosporangium haraeaeum*, *Venturia nashicola*, and *A. kikuchiana* in pears. II 500-1000, Cu hydroxide chloride 1470, and 8-hydroxyquinolinatocopper [10380-28-6] 772 µg/mL inhibited *Pseudoperonospora cubensis*, *Sphaerotheca fuliginea*, and *Pseudomonas lachrymans* in cucumber. The inhibitory mechanism of II on the growth of pathogenic bacilli includes leakage of biotin, glucose, ATP, and protein from the bacilli.

L18 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1964:496311 CAPLUS Full-text

DOCUMENT NUMBER: 61:96311

ORIGINAL REFERENCE NO.: 61:13822g-h

TITLE: Modes of action of chemotherapeutic agents in plants.

Discussion

AUTHOR(S): Cowling, Ellis B.; et al.

CORPORATE SOURCE: Conn. Agr. Expt. Sta., New Haven

SOURCE: Conn. Agr. Expt. Sta., New Haven, Bull. No. (1963), 663, 72-7

DOCUMENT TYPE: Journal

LANGUAGE: Unavailable

AB Chemical differences between pathogens and their plant hosts are considered, with some apparently new data. Relations between phenols and carbohydrate metabolism are discussed. In expts. on

fusiform rust (a major disease of southern pine trees), the stem-invading fungus produces stem galls. Cycloheximide (I) in very low concns. prevented the germination of rust spores. I was translocated in slash pine seedlings at concns. high enough to inhibit a test-assay organism (not named) but had no apparent effect on the fungus in the tissue of the infected host. It is possible that I did not diffuse to the sites of infection rapidly enough to affect the pathogen. The relative fungicidal concns. of ethionine (II) on agar (test fungus not named) were 25, 50, and over 1000 p.p.m. for the L-, DL-, and D-forms, resp. Possibly II acted as a competitive inhibitor for methionine required as a Me donor in the formation of pectin. Applications of HgCl<sub>2</sub> or CuCl<sub>2</sub> to the endocarp of pea pods induced the formation of pisatin in concns. which inhibited some pathogens of peas in vitro. Other chemical compds. induced the formation of lower concns. of pisatin.

```
=> s ?benzamide? and (anti!fung? or fungicid? or pesticid? or
herbicid?)
      35673 ?BENZAMIDE?
        3 ANTI!FUNG?
      118517 FUNGICID?
      98639 PESTICID?
      93940 HERBICID?
L19    1540 ?BENZAMIDE? AND (ANTI!FUNG? OR FUNGICID? OR PESTICID? OR
HERBICI
      D?)

=> s L19 and (py<2003 or ay<2003 or pry<2003)
      22983274 PY<2003
      4503738 AY<2003
      3972615 PRY<2003
L20    1194 L19 AND (PY<2003 OR AY<2003 OR PRY<2003)

=> s L20 and (methionine) and (spore germin?)
      97295 METHIONINE
      557 METHIONINES
      97489 METHIONINE
          (METHIONINE OR METHIONINES)
      25587 SPORE
      22994 SPORES
      38192 SPORE
          (SPORE OR SPORES)
      74630 GERMIN?
      7001 SPORE GERMIN?
          (SPORE(W)GERMIN?)
L21    0 L20 AND (METHIONINE) AND (SPORE GERMIN?)

=> s L20 and methionine
      97295 METHIONINE
      557 METHIONINES
      97489 METHIONINE
          (METHIONINE OR METHIONINES)
L22    4 L20 AND METHIONINE
```

=> d 122 ibib abs 1-4

L22 ANSWER 1 OF 4 CAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 2003:57898 CAPLUS Full-text  
DOCUMENT NUMBER: 138:122646  
TITLE: Preparation of imidazolemethanamines and  
methods for the inhibition of protozoal, fungal and/or  
bacterial agents such as Trypanosoma cruzi  
INVENTOR(S): Hamilton, Andrew D.; Van Voorhis, Wesley C.;  
Yokoyama, Kohei; Buckner, Frederick S.; Ohkanda, Junko;  
Gelb, Michael  
PATENT ASSIGNEE(S): Yale University, USA; University of Washington  
SOURCE: PCT Int. Appl., 59 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	----
-----				
WO 2003006012	A1	20030123	WO 2002-US22195	
20020711 <--				
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,				
CH, CN,				
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD,				
GE, GH,				
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,				
LK, LR,				
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ,				
OM, PH,				
PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR,				
TT, TZ,				
UA, UG, US, UZ, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT,				
BE, BG,				
CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU,				
MC, NL,				
PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,				
ML, MR,				
NE, SN, TD, TG				
CA 2453396	A1	20030123	CA 2002-2453396	
20020711 <--				
AU 2002322465	A1	20030129	AU 2002-322465	
20020711 <--				
BR 2002011098	A	20041109	BR 2002-11098	
20020711 <--				
US 20060167269	A1	20060727	US 2004-483096	
20040927 <--				
PRIORITY APPLN. INFO.:			US 2001-304711P	P
20010711 <--				

20020711 &lt;---

OTHER SOURCE(S):

MARPAT 138:122646

GI



AB The present invention relates to imidazolemethanamines (I; variables defined below; e.g. Me 2-phenyl-4-[[[1-(4-phenylbenzyl)imidazol-5-yl)methyl]amino]benzoate). For I: RA is a C1-C10 (un)substituted linear, branch-chained or cyclic alkyl or alkenyl group or a (un)substituted Ph; RB is a C1-C10 (un)substituted linear, branch-chained or cyclic alkyl or alkenyl group or a (un)substituted Ph group; and R11 and R12 = H or a C1-C3 alkyl or alkenyl group. I can be used to treat infections caused by protozoal, fungal and/or bacterial agents such as *Trypanosoma cruzi*, *Mycobacterium* spp., *Leishmania* spp., *Cryptococcus* spp., *Aspergillus* spp., *Histoplasma* spp., *Candida* spp. especially *Candida albicans*, *Pneumocystis carinii*, *Trichophyton* spp., *Microsporum* spp., *Malassezia* spp., *Rhizopus* spp., *Pseudallescheria* spp., *Blastomyces dermatitidis* and *Coccidioides* spp., among others. EC50 values are reported for about 40 I for inhibition of *T. cruzi* on 3T3 fibroblasts and for inhibition of fibroblast growth (an indication of potential toxicity). In general, hydrophobic substitution showed better activity than more polar ones and para substitution resulted in more potency than meta or ortho. The most potent compound was Me 2-phenyl-4-[[[1-(4-phenylbenzyl)imidazol-5-yl)methyl]amino]benzoate, with a remarkable activity of 500 pM; this is among the most potent known compds. against *T. cruzi* amastigotes. Even the analog without the ester group (1-(4-phenylbenzyl)-5-[[[biphenyl-3-yl)amino]methyl]imidazole) had an activity of 10 nM. The results for anti-*T. cruzi* activity in infected mice is much better for 1-(4-phenylbenzyl)-5-[[[biphenyl-3-yl)amino]methyl]imidazole than for the ester. 1-(4-Methylbenzyl)-5-[[[biphenyl-3-yl)amino]methyl]imidazole was tested for anti-*Candida* activity against a number of strains of fungus; for some strains, this compound exhibited greater activity than Fluconazole. Although the methods of preparation are not several example preps. are included and characterization data is included for about 40 I.

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS

RE FORMAT RECORD. ALL CITATIONS AVAILABLE IN THE

DOCUMENT NUMBER: 129:8597  
 ORIGINAL REFERENCE NO.: 129:1853a,1856a  
 TITLE: Embedding and encapsulation of controlled release

INVENTOR(S): particles  
 Van Lengerich, Bernhard H.  
 PATENT ASSIGNEE(S): Van Lengerich, Bernhard H., USA  
 SOURCE: PCT Int. Appl., 63 pp.  
 CODEN: PIXXD2

DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9818610	A1	19980507	WO 1997-US18984	
19971027 <--				
W: AU, CA, JP, NO, PL, US				
RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
CA 2269806	A1	19980507	CA 1997-2269806	
19971027 <--				
CA 2269806	C	20060124		
AU 9749915	A	19980522	AU 1997-49915	
19971027 <--				
AU 744156	B2	20020214		
EP 935523	A1	19990818	EP 1997-912825	
19971027 <--				
EP 935523	B1	20040929		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, FI				
JP 2002511777	T	20020416	JP 1998-520558	
19971027 <--				
EP 1342548	A1	20030910	EP 2003-10031	
19971027 <--				
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, FI				
AT 277739	T	20041015	AT 1997-912825	
19971027 <--				
PL 191399	B1	20060531	PL 1997-333095	
19971027 <--				
NO 9902036	A	19990428	NO 1999-2036	
19990428 <--				
PRIORITY APPLN. INFO.:			US 1996-29038P	P
19961028 <--				
			US 1997-52717P	P
19970716 <--			EP 1997-912825	A3
19971027 <--				
			WO 1997-US18984	W
19971027 <--				
AB				
Controlled release, discrete, solid particles which contain an encapsulated and/or embedded component such as a heat sensitive or				

readily oxidizable pharmaceutically, biol., or nutritionally active component are continuously produced without substantial destruction of the matrix material or encapsulant. A release-rate controlling component is incorporated into the matrix to control the rate of release of the encapsulant from the particles. The addnl. component may be a hydrophobic component or a high water binding capacity component for extending the release time. The plasticizable matrix material, such as starch, is admixed with at least one plasticizer, such as water, and at least one release-rate controlling component under low shear mixing conditions to plasticize the plasticizable material without substantially destroying the at least one plasticizable material and to obtain a substantially homogeneous plasticized mass. The plasticizer content is substantially reduced and the temperature of the plasticized mass is substantially reduced prior to admixing the plasticized mass with the encapsulant to avoid substantial destruction of the encapsulant and to obtain a formable, extrudable mixture. The mixture is extruded through a die without substantial or essentially no expansion and cut into discrete, relatively dense particles. Release properties may also be controlled by precoating the encapsulant and/or coating the extruded particles with a film-forming component. An example of encapsulation of acetylcysteine is given using starch, polyethylene, glycerol monostearate, and vegetable oil.

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

L22 ANSWER 3 OF 4 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1993:595101 CAPLUS Full-text

DOCUMENT NUMBER: 119:195101

ORIGINAL REFERENCE NO.: 119:34529a,34532a

TITLE: Rational estimation of the QSAR (quantitative structure-activity relationships) descriptors  $\sigma S^\circ$ , and their applications for medicinals now available

AUTHOR(S): Sasaki, Yoshio; Takagi, Tatsuya; Kawaki, Hideko

CORPORATE SOURCE: Fac. Pharm. Sci., Osaka Univ., Suita, 565, Japan

SOURCE: Chemical & Pharmaceutical Bulletin (1993), 41(3), 415-23

CODEN: CPBTAL; ISSN: 0009-2363

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Rational estimation of the descriptor  $\sigma S^\circ$  (substituent entropy constant), representing the dispersion and repulsion energies in the van der Waals interaction for both aliphatic and aromatic moieties, enabled the authors to present the descriptors of several important medicines now available. In this work, the fundamental role for the estimation of the descriptor for a substrate having a variety of binding modes and the correction value  $\Delta S_0$  necessary for aliphatic heterocycle formation are confirmed, and the descriptors for several important moieties are established according to the concept of quant. structure-activity



relationship analogy. Furthermore, several kinds of herbicides, antiinflammatory agents, hypocholesterolemic, analgesics, sympathetic stimulants, and antipsychotics are concerned in this work.

L22 ANSWER 4 OF 4 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1966:52058 CAPLUS Full-text  
DOCUMENT NUMBER: 64:52058  
ORIGINAL REFERENCE NO.: 64:9729g-h,9730a-e  
TITLE: N-Substituted derivatives of Mitomycin A and Mitomycin C  
INVENTOR(S): Meyer, Walter E.; Patrick, James B.; Mowat, John H.  
PATENT ASSIGNEE(S): American Cyanamid Co.  
SOURCE: 4 pp.  
DOCUMENT TYPE: Patent  
LANGUAGE: Unavailable  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	----

-----				
US 3226393		19651228	US 1962-200631	

19611109 <--

PRIORITY APPLN. INFO.: US

19611109 <--

GI For diagram(s), see printed CA Issue.

AB I, where X is OMe or H<sub>2</sub>N and R is alkyl or acyl, were prepared by acylation or alkylation of mitomycin A (I) (X is OMe, R is H) (II) or mitomycin C (I) (X is H<sub>2</sub>N, R is H) (III). Thus, to 41.5 mg. NaHCO<sub>3</sub> in 1.25 ml. H<sub>2</sub>O were added 1.25 ml. HCONMe<sub>2</sub> (DMF) and 10 mg. II, followed by 0.5 ml. acid-free MeI. The mixture was stirred 5 hrs. kept overnight, aerated with N, and concentrated. The residue was extracted with CHCl<sub>3</sub>, the extract concentrated, and the residue treated with ether to give N-methylmitomycin A (IV), purified by liquid-liquid partition chromatography, then crystallized from CCl<sub>4</sub> and heptane. Also, 28.8 mg. porfiromycin (I) (X is NH<sub>2</sub>, R is Me) in 5 ml. NaOH was kept 45 min. while 1 mole NH<sub>3</sub> evolved. The solution was neutralized to pH 7.0, concentrated, the residue extracted with tetrahydrofuran, the solution cooled to 5°, and excess CH<sub>2</sub>N<sub>2</sub> in ether added. The product was purified by liquid-liquid partition chromatography by using 70:30:17:4 heptane-EtOAcMeOH-H<sub>2</sub>O to give IV identical with that prepared from II. Similarly, from II were prepared N-ethylmitomycin A, N-(p-bromophenacyl)mitomycin A (V) (precipitated from ether-petr. ether), and N-benzylmitomycin A (VI). A solution of 0.02 g. II in 0.75 ml. DMF was stirred with 0.05 g. Ag<sub>2</sub>O and 0.1 ml. MeI 1 hr., then diluted with 4 vols. CHCl<sub>3</sub> to give IV. To 167 mg. carbonyldiimidazole in 2.5 ml. CHCl<sub>3</sub> was added 0.05 ml. HOAc. After 45 min. at 25°, 20 mg. II in 1 ml. CHCl<sub>3</sub> was added. After 18 hrs. the solution afforded N-acetylmitomycin A (VII), precipitated from ether with petr. ether. Similarly were prepared the p-iodobenzoyl, isonicotinoyl, and 4-iodo-3-nitrobenzoyl derivs. of II. p-Iodophenyl isocyanate (VIII) was prepared by

refluxing 200 mg. p-iodobenzamide in 9 ml. PhMe for 90-min. The solution was then diluted with 8 ml. CHCl<sub>3</sub> and added to 50 mg. II in 4 ml. CHCl<sub>3</sub>. After 24 hrs., 0.25 ml. EtOH was added; later the solution was concentrated, the residue treated (taken up in ether and the solution diluted with petr. ether) twice to get rid of Et p-iodophenylcarbamate, leaving N-(p-iodophenylcarbamoyl)mitomycin A, recrystd. from C<sub>6</sub>H<sub>6</sub>. Similarly, a solution of VIII was added to a suspension of 50 mg. III in 5 ml. CHCl<sub>3</sub> plus 0.12 ml. pyridine. After the addition of EtOH, dilution with petr. ether gave a precipitate which was taken up in EtOAc and repptd. with petr. ether to yield N-(p-iodophenylcarbamoyl)mitomycin C, recrystd. from EtOAc-petr. ether. A solution of 100 mg. II in 8 ml. CHCl<sub>3</sub> was treated with 0.45 ml. (iso-Pr)<sub>2</sub>NEt and then with 200 mg. p-BrC<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>Cl in 4 ml. CHCl<sub>3</sub>. After 24 hrs., workup afforded N-(p-bromobenzenesulfonyl)mitomycin A (IX), which was purified by partition chromatography and crystallized from CH<sub>2</sub>Cl<sub>2</sub>-C<sub>6</sub>H<sub>6</sub> as the 0.5C<sub>6</sub>H<sub>6</sub> solvate. To 0.710 g. II in 1.0 ml. CHCl<sub>3</sub> containing 0.1020 g. Et<sub>3</sub>N was added 0.0990 g. ClCO<sub>2</sub>Et in 1.0 ml. CHCl<sub>3</sub>. After 20 hrs. the mixture was worked up to give N-(carbethoxy)mitomycin A (X), m. 158-62° (purple crystals from EtOH-petr. ether) with loss of birefringence at 140°. When X was exposed to dilute acids, it was converted to a compound with an uv spectrum similar to that of apomitomycin A. To 0.025 g. VI in 0.25 ml. MeOH at 0° was added 5 ml. MeOH saturated with NH<sub>3</sub> at 0°. After storage 20 hrs. at 0° the mixture yielded N-(benzyl)mitomycin C, precipitated from CHCl<sub>3</sub>-ether with heptane. I are useful as antibacterials. Antifungal and antibacterial activity in terms of min. inhibitory concns. against 19 microorganisms is tabulated for II, V, VI, IX, X, IV, and VII. Tests in vivo showed IV was less toxic than II.

=> s 120 and (spore) and germin?

25587 SPORE

22994 SPORES

38192 SPORE

(SPORE OR SPORES)

74630 GERMIN?

L23

2 L20 AND (SPORE) AND GERMIN?

=> d 123 ibib abs 1-2

L23 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1964:19286 CAPLUS Full-text

DOCUMENT NUMBER: 60:19286

ORIGINAL REFERENCE NO.: 60:3429h,3430f

TITLE: Fungicidal activity of chloronitrobenzonitriles

AUTHOR(S): Koopmans, M. J.

CORPORATE SOURCE: N.V. Philips-Duphar, Weesp, Neth.

SOURCE: Mededelingen van de Landbouwhogeschool en de Opzoekingsstations van de Staat te Gent (1962), 27(3), 1204-13

CODEN: MLOSAT; ISSN: 0369-0695

DOCUMENT TYPE:

Journal

LANGUAGE: Dutch

AB For the assessment of the fungicidal activity a spore germination test with 3 species of fungi was used and for the assessment of phytotoxicity of some compds. the degree of leaf damage in 5 species of green plants. The activity was determined of all isomers of Cl(nO2N)mC6H5(n+m)CN, with n = 0, 1, 2, 3 and m = 0, 1, and 2, and in 21 related compds. in which the CN group had been substituted by another radical or by H. Fungal toxicity is expressed as min. lethal dose in p.p.m. The substitution of NO2 and Cl groups increases the toxicity (from >1000 p.p.m. for PhCN to 0.1 p.p.m. for 2,4,6-trichloro-3,5-dinitrobenzo-nitrile). Substitution of the CN by COOH, CONH2, CHO, CH2OH or SO2NH2 diminishes fungal toxicity considerably. The phytotoxicity of the chloronitrobenzonitriles is inversely proportional to the number of Cl atoms.

L23 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1961:114088 CAPLUS

DOCUMENT NUMBER: 55:114088

ORIGINAL REFERENCE NO.: 55:21466h-i,21467a

TITLE: Chlorocyclopentanones as nematocides and fungicides

INVENTOR(S): Richter, Sidney B.; Wahlborg, Harold J.

PATENT ASSIGNEE(S): Velsicol Chemical Corp.

DOCUMENT TYPE: Patent

LANGUAGE: Unavailable

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	----
---				
US 2980579		19610418	US	

<---

AB 2,3,4,4,5,5-Hexachloro-2-cyclopenten-1-one (I) derivs. possess fungicide and nematocide activity. I was prepared according to the method of Newcomer and McBee (CA 43, 4230e). Then 39 g. I, 69 g. AcCl, and 10 drops concentrated H2SO4, refluxed for 1 hr., allowed to stand overnight, diluted with H2O, filtered, and recrystd. from ether-heptane gave 74% yield of 3-acetylrimino-2,2,4,4,5-pentachlorocyclopentanone (II), m.p. 136-8°. Similarly, analogs of II were prepared (m.p. given): 3-acryloylimino, 129-30° (MeOH); 3-caproylimino, 62-5° (ligroine); 3-chloroacetylrimino, 121-3° (Et2O-hexane); 3-benzoylimino, 154-6° (Et2O); 3-(p-chlorobenzoylimino), 145-7° (Et2O); and 3-(o-chlorobenzoylimino), 134-6° (C6H6-hexane). These compds. at 100 p.p.m. gave inhibition of fungus spore germination, control of late blight (Phytophthora infestans) disease on foliage, and kill of the nematode Panagrellus redivivus.

=> s ?benzamide? and ?pyrimidine?

35673 ?BENZAMIDE?

95928 ?PYRIMIDINE?

L24 1640 ?BENZAMIDE? AND ?PYRIMIDINE?

=> s l24 and (anti!fung? or fungicid? or pesticid? or herbicid?)  
3 ANTI!FUNG?  
118517 FUNGICID?  
98639 PESTICID?  
93940 HERBICID?

L25 93 L24 AND (ANTI!FUNG? OR FUNGICID? OR PESTICID? OR  
HERBICID?)

=> s l25 and synerg?  
128176 SYNERG?

L26 5 L25 AND SYNERG?

=> d l26 ibib abs 1-5

L26 ANSWER 1 OF 5 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2008:1506931 CAPLUS Full-text

DOCUMENT NUMBER: 150:29914

TITLE: Pesticidal composition comprising a  
strigolactone derivative and a fungicide  
compound

INVENTOR(S): Suty-Heinze, Anne; Vors, Jean-Pierre

PATENT ASSIGNEE(S): Bayer Cropscience SA, Fr.

SOURCE: PCT Int. Appl., 41pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent


LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	----
WO 2008152092	A2	20081218	WO 2008-EP57385	
20080612				
W:	AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW,			
BY, BZ,	CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE,			
EG, ES,	FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS,			
JP, KE,	KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY,			
MA, MD,	ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,			
PG, PH,	PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY,			
TJ, TM,	TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW			
HR, HU,	RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR,			
SI, SK,	IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE,			
SN, TD,	TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE,			
ZM, ZW,	TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG,			

AM, AZ, BY, KG, KZ, MD, RU, TJ, TM  
PRIORITY APPLN. INFO.: EP 2007-356084 A  
20070615  
OTHER SOURCE(S): MARPAT 150:29914  
GI



AB The invention relates to a pesticidal composition comprising a strigolactone derivative (a) and a fungicide compound (b) in a weight ratio of (a)/(b) ranging from 1/1 to 1/1014; such a composition may include an addnl. fungicidal compound and may be supplemented with arbuscular mycorrhizal fungi. A method for preventively or curatively controlling phytopathogenic fungi of crops with a composition according to the invention and use of this composition to control phytopathogenic fungi and parasitic weed species are claimed also. In a microtest performed with *Pyricularia oryzae*, a synergistic effect in controlling fungal growth was found with the mixture of trifloxystrobin 0.3 + I 0.00003 ppm.

L26 ANSWER 2 OF 5 CAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 2006:512967 CAPLUS Full-text  
DOCUMENT NUMBER: 144:482751  
TITLE: Synergistic fungicidal menadione compositions  
INVENTOR(S): Koehle, Harald; Stierl, Reinhard; Gold,  
Randall Evan; Goerth, Felix Christian; Speakman, John-Bryan;  
Dombo, Peter; Semar, Martin; Strobel, Dieter;  
Niedenbrueck, Matthias; Bestman, Hans  
PATENT AGENT(S): Basf Aktiengesellschaft, Germany  
SOURCE: PCT Int. Appl., 43 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: German  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2006056434	A1	20060601	WO 2005-EP12562	

20051124  
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,  
CA, CH,  
CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,  
GB, GD,  
GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN,  
KP, KR,  
KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN,  
MW, MX,  
MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,  
SD, SE,  
SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US,  
UZ, VC,  
VN, YU, ZA, ZM, ZW  
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR,  
HU, IE,  
IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR,  
BF, BJ,  
CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG,  
BW, GH,  
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM,  
AZ, BY,  
KG, KZ, MD, RU, TJ, TM  
EP 1819223 A1 20070822 EP 2005-809496  
20051124  
EP 1819223 B1 20080312  
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR,  
HU, IE,  
IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK,  
TR, HR, YU  
AT 388635 T 20080315 AT 2005-809496  
20051124  
BR 2005017881 A 20081021 BR 2005-17881  
20051124  
US 20080039320 A1 20080214 US 2007-791464  
20070523  
PRIORITY APPLN. INFO.: DE 2004-102004057279A  
20041126  
WO 2005-EP12562 W

20051124  
OTHER SOURCE(S): MARPAT 144:482751  
AB Synergistic fungicidal compns. comprise menadione and at least one agent selected from: (A) azoles, such as cyproconazole, difenoconazole, epoxiconazole, fluquinconazole, flusilazole, hexaconazole, imazalil, metconazole, myclobutanil, penconazole, prochloraz, prothioconazole, tebuconazole, triadimefon, triadimenol, triflumizole; (B) strobilurines, such as azoxystrobin, dimoxystrobin, fluoxastrobin, kresoxim-Me, metominostrobin, orysastrobin, picoxystrobin, pyraclostrobin, or trifloxystrobin; (C) acylalanines, such as benalaxyl, metalaxyl, mefenoxam, ofurace, oxadixyl; (D) amine derivs., such as spiroxamine; (E) anilinopyrimidines, such as pyrimethanil, mepanipyrim, or cyprodinil; (F) dicarboximides, such as iprodion, procymidon, vinclozolin; (G) cinnamides and analogs, such as dimethomorph, flumetover, or flumorph; (H) dithiocarbamates, such as ferbam, nabam, maneb, metam, metiram, propineb, polycarbamate, thiram, ziram, zineb; (I) heterocyclic

comps., such as benomyl, boscalid, carbendazim, dithianon, famoxadone, fenamidone, picobenzamide, proquinazid, quinoxifen, thiophanat-Me, triforine, 5-chloro-7-(4-methyl-piperidine-1-yl)-6-(2,4,6-trifluoro-phenyl)-[1,2,4]triazolo[1,5-a]pyrimidin, 3-(3-bromo-6-fluoro-2-methyl-indol-1-sulfonyl)-[1,2,4]triazol-1-sulfonic acid di-Me amide, or thiophene derivs.

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L26 ANSWER 3 OF 5 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2005:1262708 CAPLUS Full-text

DOCUMENT NUMBER: 143:473909

TITLE: Synergistic fungicide mixture comprising a triazolopyrimidine and a pyridine derivative

INVENTOR(S): Tormo I Blasco, Jordi; Grote, Thomas; Scherer, Maria;

Schoefl, Stierl, Reinhard; Strathmann, Siegfried;

Ulrich; Gewehr, Markus

PATENT ASSIGNEE(S): BASF Aktiengesellschaft, Germany

SOURCE: PCT Int. Appl., 18 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
2005112643	A1	20051201	WO 2005-EP4482	
20050427				
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			

20050427	AU 2005245261	A1	20051201	AU 2005-245261
20050427	CA 2562637	A1	20051201	CA 2005-2562637
20050427	EP 1748692	A1	20070207	EP 2005-742678
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, LV				
20050427	CN 1949973	A	20070418	CN 2005-80014599
20050427	BR 2005010489	A	20071113	BR 2005-10489
20050427	JP 2007536305	T	20071213	JP 2007-511955
20050427	MX 2006011749	A	20070116	MX 2006-11749
20061011	IN 2006KN02974	A	20070608	IN 2006-KN2974
20061013	US 20070191398	A1	20070816	US 2006-579672
20061107	NO 2006005508	A	20061201	NO 2006-5508
20061129	KR 2007011576	A	20070124	KR 2006-725650
20061206	PRIORITY APPLN. INFO.:			DE 2004-102004023248A
20040507				WO 2005-EP4482 W

20050427

AB A synergistic fungicide mixture comprises 5-chloro-7-(4-methylpiperidin-1-yl)-6-(2,4,6-trifluorophenyl)-[1,2,4]triazolo[1,5-a]pyrimidine and 2,6-dichloro-N-(3-chloro-5-trifluoromethylpyridin-2-ylmethyl) benzamide.

REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L26 ANSWER 4 OF 5 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2005:1106849 CAPLUS Full-text

DOCUMENT NUMBER: 143:361642

TITLE: Synergistic ternary fungicidal mixtures

INVENTOR(S): Tormo i Blasco, Jordi; Grote, Thomas; Scherer, Maria;

Schoefl, Stierl, Reinhard; Strathmann, Siegfried;

Ulrich

PATENT ASSIGNEE(S): BASF Aktiengesellschaft, Germany

SOURCE: PCT Int. Appl., 38 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 6

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
------------	------	------	-----------------	------



-----	-----	-----	-----	-----
WO 2005094583	A1	20051013	WO 2005-EP3213	
20050326				
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,				
CA, CH,				
CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,				
GB, GD,				
GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,				
KZ, LC,				
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,				
NA, NI,				
NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK,				
SL, SM,				
SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU,				
ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,				
ZW, AM,				
AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ,				
DE, DK,				
EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL,				
PL, PT,				
RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ,				
GW, ML,				
MR, NE, SN, TD, TG				
AU 2005227688	A1	20051013	AU 2005-227688	
20050326				
CA 2558062	A1	20051013	CA 2005-2558062	
20050326				
EP 1732388	A1	20061220	EP 2005-729121	
20050326				
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR,				
HU, IE,				
IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR,				
HR, LV, YU				
CN 1937920	A	20070328	CN 2005-80010641	
20050326				
BR 2005008965	A	20070821	BR 2005-8965	
20050326				
JP 2007537156	T	20071220	JP 2007-505466	
20050326				
IN 2006KN02365	A	20070525	IN 2006-KN2365	
20060821				
MX 2006009693	A	20061116	MX 2006-9693	
20060825				
NO 2006004923	A	20061027	NO 2006-4923	
20061027				
KR 2007004068	A	20070105	KR 2006-722407	
20061027				
PRIORITY APPLN. INFO.:			DE 2004-102004016084A	
20040330				
			WO 2005-EP3213	W
20050326				
AB	Synergistic ternary fungicidal mixts. comprise 5-chloro-7-(4-methylpiperidin-1-yl)-6-(2,4,6-trifluorophenyl)-[1,2,4]triazolo[1,5-a]pyrimidine, a strobilurin derivative (pyraclostrobin or orysastrobin) and a fungicide selected from			

acylalanines, amine derivs., anilinoypyrimidines, antibiotics, azoles, dicarboximides, dithiocarbamates, copper fungicides, nitrophenyl derivs., phenylpyrroles, sulfenic acid derivs., cinnamic acid derivs. and their analogs and anilazine, benomyl, boscalid, carbendazim, carboxin, oxycarboxin, cyazofamid, dazomet, dithianon, famoxadone, fenamidone, fenarimol, fuberidazole, flutolanil, furametpyr, isoprothiolane, mepronil, nuarimol, picobenzamide, probenazole, proquinazid, pyrifenoxy, pyroquilon, quinoxifen, silthiofam, thiabendazole, thifluzamide, thiophanate-Me, tiadinil, tricyclazole, triforine, sulfur, acibenzolar-S-Me, benthiavalicarb, carpropamid, chlorothalonil, cyflufenamid, cymoxanil, dazomet, diclomezine, diclocymet, diethofencarb, edifenphos, ethaboxam, fenhexamid, fentin acetate, fenoxanil, ferimzone, fluazinam, phosphorous acid, fosetyl, fosetyl-aluminum, iprovalicarb, hexachlorobenzene, metrafenone, penicuron, propamocarb, phthalide, tolclofos-Me, quintozene and zoxamideamt.

REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L26 ANSWER 5 OF 5 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2004:550533 CAPLUS Full-text

DOCUMENT NUMBER: 141:82297

TITLE: Immunostimulatory nucleic acids for the treatment of

disorders associated with microorganisms, for preventing antibiotic resistance and for

treating and

preventing warts

INVENTOR(S): Bratzler, Robert L.; Petersen, Deanna M.

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 54 pp., Cont. of U.S. Ser. No.

801,839, abandoned.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	----
-----				
US 20040131628	A1	20040708	US 2003-666733	
20030919				
PRIORITY APPLN. INFO.:			US 2000-187834P	P
20000308				
			US 2001-801839	B1

20010308

OTHER SOURCE(S): MARPAT 141:82297

AB The invention involves administration of an immunostimulatory nucleic acid alone or in combination with an antimicrobial agent for the treatment or prevention of infectious disease associated with microorganisms in subjects, for preventing antibiotic resistance and for treating and preventing warts. The combination of drugs are administered in synergistic amts. or in various

dosages or at various time schedules. The invention also relates to kits and compns. concerning the combination of drugs.

=> s ?benzamide? and ?pyrimidine?

35673 ?BENZAMIDE?

95928 ?PYRIMIDINE?

L27 1640 ?BENZAMIDE? AND ?PYRIMIDINE?

=> s L27 and (?carboxamide? or phthalamid?)

44983 ?CARBOXAMIDE?

1128 PHTHALAMID?

L28 506 L27 AND (?CARBOXAMIDE? OR PHTHALAMID?)

=> s L28 and (mycelium)

16266 MYCELIUM

29 MYCELIUMS

9113 MYCELIA

2 MYCELIAS

23446 MYCELIUM

(MYCELIUM OR MYCELIUMS OR MYCELIA OR MYCELIAS)

L29 0 L28 AND (MYCELIUM)

=> s L28 and spor? and germinat?

90964 SPOR?

63685 GERMINAT?

L30 0 L28 AND SPOR? AND GERMINAT?

=> s L28 and methionine

97295 METHIONINE

557 METHIONINES

97489 METHIONINE

(METHIONINE OR METHIONINES)

L31 8 L28 AND METHIONINE

=> d L31 ibib abs 1-8

L31 ANSWER 1 OF 8 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2007:912269 CAPLUS Full-text

DOCUMENT NUMBER: 147:277915

TITLE: Preparation of 4-phenylpiperidine-substituted amino

acid derivatives, particularly valine amides, as

modulators of chemokine receptor activity and their

use in the treatment of inflammatory and autoimmune

diseases



INVENTOR(S): Carter, Percy H.; Cavallaro, Cullen L.; Duncia, John

V.; Gardner, Daniel S.; Hynes, John; Liu, Rui-Qin;

Santella, Joseph B.; Dodd, Dharmpal S. PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA

SOURCE: PCT Int. Appl., 515pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
20070125	A2	20070816	WO 2007-US61012	
20070125				
CA, CH,				
GB, GD,				
KM, KN,				
MG, MK,				
PT, RO,				
TR, TT,				
HU, IE,				
BF, BJ,				
BW, GH,				
AZ, BY,				
US 20070208056	A1	20070906	US 2007-625874	
20070123				
AU 2007212236	A1	20070816	AU 2007-212236	
20070125				
IN 2008DN06339	A	20081024	IN 2008-DN6339	
20080721				
KR 2008095890	A	20081029	KR 2008-720904	
20080826				
PRIORITY APPLN. INFO.:				
20060127				
20070123				
20070125				
OTHER SOURCE(S):				
GI				



AB Title compds. I [T = CO, COO, CONH, CON-alkyl, SO<sub>2</sub>; R<sub>1</sub> = (un)substituted cyclo/alkyl, (hetero)aryl, heterocyclyl; R<sub>2</sub> = cycloalkyl/cyclo/alkyl, alkenyl optionally substituted with OH; R<sub>3</sub> at each occurrence = alkyl; or any 2 R<sub>3</sub>'s attached to the same C may form a 3-6 membered ring; W = H, F, OH, CN, NH<sub>2</sub>; R<sub>5</sub> = halo, CN, alkoxy; W and one R<sub>5</sub> together with the C atoms to which each is attached may form an (un)substituted 3-6 membered O containing ring; m at each occurrence = independently 0-2; n = 1-3; and their stereoisomers, prodrugs and pharmaceutically acceptable salts] were prepared as modulators of CCR-1 and MIP-1, especially MIP-1α receptors. Thus, valine amide II was prepared using N-(tert-butoxycarbonyl)-D-valine, 4-(4-chlorophenyl)piperidine hydrochloride, and benzoic acid. All the invention compds. were evaluated for their chemokine receptor modulatory activity. Methods of treating and preventing inflammatory diseases such as asthma and allergic diseases, as well as autoimmune pathologies such as rheumatoid arthritis and atherosclerosis using said modulators are disclosed.

L31 ANSWER 2 OF 8 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2007:646507 CAPLUS [Full-text](#)

DOCUMENT NUMBER: 147:249819

TITLE: Development of Reliable Aqueous Solubility Models and

Their Application in Druglike Analysis

AUTHOR(S): Wang, Junmei; Krudy, George; Hou, Tingjun; Zhang, Wei;

Holland, George; Xu, Xiaojie

CORPORATE SOURCE: Encysive Pharmaceuticals Inc., Houston, TX, 77030, USA

SOURCE: Journal of Chemical Information and Modeling (2007),

47(4), 1395-1404

CODEN: JCISD8; ISSN: 1549-9596

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB In this work, two reliable aqueous solubility models, ASMS (aqueous solubility based on mol. surface) and ASMS-LOGP (aqueous solubility based on mol. surface using calculated log P (ClogP) as a descriptor), were constructed by using atom type classified solvent accessible surface areas and several mol. descriptors for a diverse data set of 1708 mols. For ASMS (without using ClogP as a descriptor), the leave-one-out q<sub>2</sub> and root-mean-square error (RMSE) were 0.872 and 0.748 log unit, resp. ASMS-LOGP was slightly better than ASMS (q<sub>2</sub> = 0.886, RMSE = 0.705). Both models were extensively validated by three cross-validation tests and encouraging predictability was achieved. High throughput aqueous solubility prediction was conducted for a number of data sets extracted from several widely used databases. The authors found that real drugs are about 20-fold more soluble than the so-called druglike mols. in the ZINC database, which have no violation of Lipinski's "Rule of 5" at all. Specifically, oral drugs are about 16-fold more soluble, while injection drugs are 50-60-fold more soluble. If the criterion of a mol. to be soluble is set to -5 log unit, about 85% of real drugs are predicted as soluble; in contrast only 50% of druglike mols. in ZINC are soluble. The authors concluded that the two models could be served as a rule in druglike anal. and an efficient filter in prioritizing compound libraries prior to high throughput screenings (HTS).

REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

L31 ANSWER 3 OF 8 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2007:593348 CAPLUS Full-text

DOCUMENT NUMBER: 147:31090

TITLE: Oxazolidinonecarboxamides as inhibitors of diacylglycerol acyltransferase (DGAT) and

their preparation, pharmaceutical compositions and

use in the treatment of obesity, diabetes type II and metabolic syndrome

INVENTOR(S): Bolin, David Robert; Cheung, Adrian Wai-Hing; Firooznia, Fariborz; Hamilton, Matthew

Michael; Li, Shiming; McDermott, Lee Apostle; Qian, Yimin;

Yun, Weiya

PATENT ASSIGNEE(S): F. Hoffmann-La Roche A.-G., Switz.

SOURCE: PCT Int. Appl., 201pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent


LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
----	----	-----	-----	----
WO 2007060140	A2	20070531	WO 2006-EP68611	

20061117  
 WO 2007060140 A3 20070913  
 W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,  
 CA, CH,  
 CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,  
 GB, GD,  
 GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG,  
 KM, KN,  
 KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD,  
 MG, MK,  
 MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL,  
 PT, RO,  
 RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN,  
 TR, TT,  
 TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW  
 RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR,  
 HU, IE,  
 IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR,  
 BF, BJ,  
 CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG,  
 BW, GH,  
 GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM,  
 AZ, BY,  
 KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA  
 AU 2006316560 A1 20070531 AU 2006-316560  
 20061117  
 CA 2630269 A1 20070531 CA 2006-2630269  
 20061117  
 US 20070123504 A1 20070531 US 2006-601429  
 20061117  
 EP 1963313 A2 20080903 EP 2006-830027  
 20061117  
 R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR,  
 HU, IE,  
 IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR  
 IN 2008DN04199 A 20080801 IN 2008-DN4199  
 20080516  
 MX 200806568 A 20080530 MX 2008-6568  
 20080521  
 KR 2008063865 A 20080707 KR 2008-712699  
 20080527  
 CN 101316844 A 20081203 CN 2006-80044426  
 20080528  
 PRIORITY APPLN. INFO.: US 2005-740578P P  
 20051128  
 US 2006-849352P P  
 20061004  
 WO 2006-EP68611 W  
 20061117  
 OTHER SOURCE(S): MARPAT 147:31090  
 GI



AB Provided herein are compds. of the formula I, as well as pharmaceutically acceptable salts thereof. Compds. of formula I wherein R1 is (un)substituted aryl; R2 is C and N; R3 and R4 are independently C, N, S, and O; R5 is C, N and S; R6 is H, (halo)alkyl, halo, thioalkyl and absent; R7 is substituted pyrimidinyl, substituted pyridinyl, substituted pyrazinyl, and substituted thiazolyl; dashed lines are optional double bonds; and their pharmaceutically acceptable salts thereof, are claimed. These compds., and the pharmaceutical compns. containing them, are useful for the treatment of diseases such as, for example, obesity, type II diabetes mellitus and metabolic syndrome. Example compound II was prepared by amidation of 2-phenyl-5-trifluoromethylthiazole-4-carboxylic acid with 6-(morpholin-4-yl)pyridin-3-ylamine. All the invention compds. were evaluated for their DGAT inhibitory activity. From the assay, it was determined that compound II exhibited an IC50 value of < 0.75  $\mu$ M.

L31 ANSWER 4 OF 8 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 2004:20322 CAPLUS Full-text  
 DOCUMENT NUMBER: 140:87658  
 TITLE: Peptidomimetic modulators of cell adhesion  
 INVENTOR(S): Gour, Barbara J.; Blaschuk, Orest W.; Ali, Anmar; Ni, Feng; Chen, Zhigang; Michaud, Stephanie  
 Denise; Wang, Shaomeng; Hu, Zengjian  
 PATENT ASSIGNEE(S): Can.  
 SOURCE: U.S. Pat. Appl. Publ., 280 pp., Cont.-in-part of U.S. Ser. No. 6,982.  
 CODEN: USXXCO  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 15  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20040006011	A1	20040108	US 2003-425557	
20030428				
US 6031072	A	20000229	US 1997-893534	
19970711				
US 6326352	B1	20011204	US 2000-507102	
20000217				



US 20020168761	A1	20021114	US 2001-769145	
20010124				
US 20020151475	A1	20021017	US 2001-6982	
20011204				
US 6914044	B2	20050705		
PRIORITY APPLN. INFO.:			US 1996-21612P	P
19960712				
19970711			US 1997-893534	A1
			US 2000-491078	B2
20000124				
			US 2000-507102	A1
20000217				
			US 2001-769145	B2
20010124				
			US 2001-6982	A2

20011204

OTHER SOURCE(S): MARPAT 140:87658

AB Peptidomimetics of cyclic peptides, and compns. comprising such peptidomimetics are provided. The peptidomimetics have a three-dimensional structure that is substantially similar to a three-dimensional structure of a cyclic peptide that comprises a cadherin cell adhesion recognition sequence HAV. Methods for using such peptidomimetics for modulating cadherin-mediated cell adhesion in a variety of contexts are also provided.

L31 ANSWER 5 OF 8 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2003:154399 CAPLUS Full-text

DOCUMENT NUMBER: 138:204936

TITLE: Preparation of heterocyclic compounds as integrase

INVENTOR(S): inhibiting antiviral agents  
Kiyama, Ryuichi; Kanda, Yasuhiko; Tada, Yukio;  
Fujishita, Toshio; Kawasuji, Takashi; Takechi,

Shozo;

Fuji, Masahiro  
PATENT ASSIGNEE(S): Shionogi & Co., Ltd., Japan

SOURCE: PCT Int. Appl., 663 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
-----	WO 2003016275	A1	20030227	WO 2002-JP8108	
20020808					
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,				
CH, CN,	CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD,				
GE, GH,	GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK,				
LR, LS,	LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM,				

PH, PL,  
 TZ, UA,  
 BE, BG,  
 MC, NL,  
 ML, MR,  
 PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT,  
 UG, US, UZ, VC, VN, YU, ZA, ZM, ZW  
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT,  
 CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU,  
 PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,  
 NE, SN, TD, TG  
 CA 2452769 A1 20030227 CA 2002-2452769  
 20020808 AU 2002320703 A1 20030303 AU 2002-320703  
 20020808 EP 1422218 A1 20040526 EP 2002-749384  
 20020808 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,  
 MC, PT,  
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK  
 BR 2002011750 A 20041013 BR 2002-11750  
 20020808 CN 1558898 A 20041229 CN 2002-819869  
 20020808 MX 2004000646 A 20040318 MX 2004-646  
 20040121 US 20040229909 A1 20041118 US 2004-485394  
 20040130  
 PRIORITY APPLN. INFO.: JP 2001-245071 A  
 20010810 JP 2001-370860 A  
 20011205 JP 2002-191483 A  
 20020628 WO 2002-JP8108 W  
 20020808  
 OTHER SOURCE(S): MARPAT 138:204936  
 AB The title compds. RDC(:Z)C(Y):CRCRA [RC and RD in combination form  
 a ring with the adjacent carbon atoms, provided that the ring may  
 be a fused ring; Y represents hydroxy, mercapto, or amino; Z  
 represents oxygen, sulfur, or NH; and RA represents N-containing  
 aromatic heterocycle, etc.] are prepared Compds. of this  
 invention in vitro showed IC50 values of 0.12 µg/mL to 2.9 µg/mL  
 against integrase. Formulations are given.  
 REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE  
 FOR THIS  
 REFORMAT RECORD. ALL CITATIONS AVAILABLE IN THE

L31 ANSWER 6 OF 8 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 2002:869496 CAPLUS Full-text  
 DOCUMENT NUMBER: 137:363033  
 TITLE: Peptidomimetic modulators of cell adhesion  
 INVENTOR(S): Gour, Barbara J.; Blaschuk, Orest W.; Ali,  
 Annmar; Ni,  
 Feng; Chen, Zhigang; Michaud, Stephanie D.;  
 Wang,

PATENT ASSIGNEE(S): Shoameng; Hu, Zenjian  
 SOURCE: Can.  
 of U.S. U.S. Pat. Appl. Publ., 309 pp., Cont.-in-part  
 Ser. No. 491,078.  
 CODEN: USXXCO  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 15  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	----
20020168761	A1	20021114	US 2001-769145	
20010124				
20040058864	A1	20040325	US 2003-412701	
20030410				
US 7268115	B2	20070911		
US 20040006011	A1	20040108	US 2003-425557	
20030428				
US 20080081831	A1	20080403	US 2007-762015	
20070612				
US 7446120	B2	20081104		
PRIORITY APPLN. INFO.:			US 2000-491078	A2
20000124				
19960712			US 1996-21612P	P
19970711			US 1997-893534	A1
20000217			US 2000-507102	A1
20010124			US 2001-769145	B1
20011204			US 2001-6982	A2
20030410			US 2003-412701	A1

OTHER SOURCE(S): MARPAT 137:363033  
 AB Peptidomimetics of cyclic peptides, and compns. comprising such peptidomimetics are provided. The peptidomimetics have a three-dimensional structure that is substantially similar to a three-dimensional structure of a cyclic peptide that comprises a cadherin cell adhesion recognition sequence HAV. Methods for using such peptidomimetics for modulating cadherin-mediated cell adhesion in a variety of contexts are also provided.

L31 ANSWER 7 OF 8 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 1992:400277 CAPLUS Full-text  
 DOCUMENT NUMBER: 117:277  
 ORIGINAL REFERENCE NO.: 117:43a,46a  
 TITLE: Mechanism of allergic cross-reactions. I.  
 Multispecific binding of ligands to a mouse  
 monoclonal  
 anti-DNP IgE antibody  
 AUTHOR(S): Varga, Janos M.; Kalchschmid, Gertrud; Klein,

Georg

CORPORATE SOURCE: F.; Fritsch, Peter  
6020, Dep. Dermatol., Univ. Innsbruck, Innsbruck,

Austria

SOURCE: Molecular Immunology (1991), 28(6), 641-54  
CODEN: MOIMD5; ISSN: 0161-5890

DOCUMENT TYPE: Journal

LANGUAGE: English

AB A recently developed solid-phase binding assay was used to investigate the specificity of ligand binding to a mouse monoclonal anti-dinitrophenyl IgE (I). All DNP-amino acids, that were tested inhibited the binding of the radio-labeled I to DNP covalently attached to polystyrene microplates; however, the concentration for 50% inhibition varied within four orders of magnitude, DNP-L-serine being the most and DNP-L-proline the least potent inhibitor. In addition to DNP analogs, a large number of drugs and other compds. were tested for their ability to compete with DNP for the binding site of I. At the concentration used for screening, 59% of compds. had no significant inhibition; 19% inhibited the binding of I more than 50%. Several families of compds. (tetracyclines, polymyxins, phenothiazines, salicylates, and quinones) that were effective competitors were found. Within these families, changes in the functional groups attached to the family stem had major effects on the affinity of ligand binding. The occurrence frequencies of interactions of ligands with I is in good agreement with the semi-empirical model for multispecific antibody-ligand interactions.

L31 ANSWER 8 OF 8 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1964:34166 CAPLUS Full-text

DOCUMENT NUMBER: 60:34166

ORIGINAL REFERENCE NO.: 60:6111b,6112a-b

TITLE: Oral antidiabetics

AUTHOR(S): Budesinsky, Z.; Zikmund, E.

SOURCE: Pharmacotherapeutica, 1950-1959 (1963) 31-48  
CODEN: 13KGA8


DOCUMENT TYPE: Journal

LANGUAGE: English

AB cf. CA 54, 6563f. A review of earlier work on 3 types of compds. and the preparation and testing of 1-arylsulfonyl-5-alkylglycocyamidines and hydantoins. The preps. were made by reactions like the following:  $\text{MeC}_6\text{H}_4\text{SO}_2\text{Cl} + \text{Ca}(\text{NHCN})_2 + \text{NaOH} \rightarrow \text{MeC}_6\text{H}_4\text{SO}_2\text{N}(\text{Na})\text{CN}$  (I);  $\text{I} + \text{BrCH}_2\text{CO}_2\text{Et} \rightarrow \text{MeC}_6\text{H}_4\text{SO}_2\text{N}(\text{CN})\text{CH}_2\text{CO}_2\text{Et}$  (II);  $\text{II} + \text{RNH} \rightarrow \text{MeC}_6\text{H}_4\text{SO}_2\text{R}$  (III);  $\text{III} + \text{H}^+ \rightarrow \text{MeC}_6\text{H}_4\text{SO}_2\text{R}^+$ , where R is a 3-substituted 2-imino-4-oxo-1-imidazolidinyl group and R' is the 2-oxo analog. A similar series of chloro compds. was prepared by starting with  $\text{ClC}_6\text{H}_4\text{SO}_2\text{Cl}$ . The hypoglycemic activity of 35 such compds. is reported. After comparison of these compds. with those in the earlier studies,  $\text{MeC}_6\text{H}_4\text{SO}_2\text{N}(\text{Bu})(\text{CH}_2\text{COOH})$  (IV) was chosen for clin. trials. Thorough testing on rats showed its hypoglycemic effect to be 60-70% of that of tolbutamide (V), with a slower onset. The effect on dogs was similar but lasted longer. No chronic toxicity was found in rats given 3 times the optimal dose for a year. From trials in 3 clinics, IV was found to be of


<http://www.cas.org/support/stngen/stndoc/properties.html>

Uploading C:\Program Files\Stnexp\Queries\10588532\_2.str



```
Match level :
1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:CLASS 8:CLASS
9:CLASS 10:CLASS 11:CLASS 12:Atom 13:Atom 14:Atom 15:Atom
16:Atom 17:Atom 18:CLASS 19:CLASS 20:CLASS
```

```
=> d 132
L32 HAS NO ANSWERS
L32 STR
```



Structure attributes must be viewed using STN Express query preparation.

```
=> s sss sam
ENTER LOGIC EXPRESSION, QUERY NAME, OR (END):L32
SAMPLE SEARCH INITIATED 15:50:29 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED -          5 TO ITERATE
```

```
100.0% PROCESSED          5 ITERATIONS          0
ANSWERS
SEARCH TIME: 00.00.01
```

```
FULL FILE PROJECTIONS:  ONLINE  **COMPLETE**
                        BATCH   **COMPLETE**
PROJECTED ITERATIONS:   5 TO    234
PROJECTED ANSWERS:      0 TO    0
```

L33 0 SEA SSS SAM L32

```
=> s l32 sss full
FULL SEARCH INITIATED 15:50:34 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED -          63 TO ITERATE
```

```
100.0% PROCESSED          63 ITERATIONS          0
ANSWERS
SEARCH TIME: 00.00.01
```

L34 0 SEA SSS FUL L32

```
=>
Connecting via Winsock to STN
```

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate substance identification.

```
=> e grosjean!courneyor marie?/au
E1          2      GROSJEAN YOLAND/AU
E2          1      GROSJEAN YVES/AU
```

```

E3      0 --> GROSJEAN!COURNEYOR MARIE?/AU
E4      1      GROSJLOS JACKIE/AU
E5      1      GROSJOE L/AU
E6      1      GROSKA D E/AU
E7      1      GROSKA E/AU
E8      1      GROSKA JUDIT/AU
E9      5      GROSKA LASZLO/AU
E10     1      GROSKAMP HANS/AU
E11     7      GROSKAUFMANIS A/AU
E12     1      GROSKAUFMANIS E/AU

```

=> e grosjean-courneyor marie?/au

```

E1      2      GROSJEAN YOLAND/AU
E2      1      GROSJEAN YVES/AU
E3      0 --> GROSJEAN-COURNEYOR MARIE?/AU
E4      1      GROSJLOS JACKIE/AU
E5      1      GROSJOE L/AU
E6      1      GROSKA D E/AU
E7      1      GROSKA E/AU
E8      1      GROSKA JUDIT/AU
E9      5      GROSKA LASZLO/AU
E10     1      GROSKAMP HANS/AU
E11     7      GROSKAUFMANIS A/AU
E12     1      GROSKAUFMANIS E/AU

```

=> e grosjean marie?/au

```

E1      2      GROSJEAN MARIE EVE/AU
E2      2      GROSJEAN MARIE HELENE/AU
E3      0 --> GROSJEAN MARIE?/AU
E4      5      GROSJEAN MARTIN/AU
E5      1      GROSJEAN MAURICE B/AU
E6      1      GROSJEAN MICHEL/AU
E7      2      GROSJEAN MIREILLE/AU
E8      5      GROSJEAN N/AU
E9      1      GROSJEAN NATHALIE/AU
E10     1      GROSJEAN NICOLE/AU
E11     1      GROSJEAN NOELLE/AU
E12     5      GROSJEAN O/AU

```

=> e gouot jean?/au

```

E1      1      GOUOT JEAN M/AU
E2      21     GOUOT JEAN MARIE/AU
E3      0 --> GOUOT JEAN?/AU
E4      1      GOUOUDKOV A V/AU
E5      1      GOUPAL D P/AU
E6      1      GOUPALE D C/AU
E7      32     GOUPALOV S V/AU
E8      1      GOUPALOV SERGUEI/AU
E9      3      GOUPALOV SERGUEI V/AU
E10     1      GOUPÉE ANDREW J/AU
E11     1      GOUELL JOHNATHAN E/AU
E12     3      GOUELL M/AU

```

=> s e2

```
L35      21 "GOUOT JEAN MARIE"/AU
```

=> s l35 and (py<2003 or pry<2003 or ay,2003)


22983274 PY<2003  
 3972615 PRY<2003  
 3351 AY  
 37 AYS  
 3388 AY  
 (AY OR AYS)  
 42924 2003  
 0 AY,2003  
 (AY(W)2003)  
 L36 8 L35 AND (PY<2003 OR PRY<2003 OR AY,2003)  
 => s 135 and (py<2003 or pry<2003 or ay<2003)  
 22983274 PY<2003  
 3972615 PRY<2003  
 4503738 AY<2003  
 L37 8 L35 AND (PY<2003 OR PRY<2003 OR AY<2003)  
 => d 137 ibib abs 1-8

L37 ANSWER 1 OF 8 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 2000:209832 CAPLUS Full-text  
 DOCUMENT NUMBER: 132:218333  
 TITLE: Synergistic fungicidal compositions  
 INVENTOR(S): Chazalet, Maurice; Duvert, Patrice; Gouet,  
 Jean-Marie; Mercer, Richard  
 PATENT ASSIGNEE(S): Aventis CropScience SA, Fr.  
 SOURCE: PCT Int. Appl., 36 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: French  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000016629	A1	20000330	WO 1999-FR2223	
19990920 <--				
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN,				
CR, CU,				
CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,				
ID, IL,				
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,				
LV, MD,				
MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG,				
SI, SK,				
SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH,				
CY, DE,				
DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF,				
BJ, CF,				
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
FR 2783401	A1	20000324	FR 1998-11895	
19980921 <--				
FR 2783401	B1	20001020		
CA 2344218	A1	20000330	CA 1999-2344218	
19990920 <--				



AU 9956299	A	20000410	AU 1999-56299
19990920 <--			
EP 1115288	A1	20010718	EP 1999-943000
19990920 <--			
EP 1115288	B1	20030115	
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,			
MC, PT,			
IE, SI, LT, LV, FI, RO			
TR 200100830	T2	20010821	TR 2001-830
19990920 <--			
BR 9914179	A	20011030	BR 1999-14179
19990920 <--			
HU 2001003881	A2	20020328	HU 2001-3881
19990920 <--			
HU 2001003881	A3	20020828	
JP 2002526429	T	20020820	JP 2000-573600
19990920 <--			
AT 230927	T	20030215	AT 1999-943000
19990920 <--			
ES 2186401	T3	20030501	ES 1999-943000
19990920 <--			
NZ 510306	A	20030829	NZ 1999-510306
19990920 <--			
CN 1212767	C	20050803	CN 1999-812006
19990920 <--			
IL 142043	A	20051120	IL 1999-142043
19990920 <--			
ZA 2001002126	A	20020614	ZA 2001-2126
20010314 <--			
MX 2001002938	A	20020311	MX 2001-2938
20010320 <--			
BG 105392	A	20011031	BG 2001-105392
20010329 <--			
BG 64721	B1	20060131	
US 6753339	B1	20040622	US 2001-787631
20010518 <--			
PRIORITY APPLN. INFO.:			FR 1998-11895 A
19980921 <--			
			WO 1999-FR2223 W
19990920 <--			
OTHER SOURCE(S):	MARPAT 132:218333		
GI			



AB The title compns. comprise an imidazolinone derivative I (M = O or S; Y = F, Cl or Me; n = 0 or 1) and ROCONHCHR1CONHCHMeA (II) [R, R1 = alkyl; A = (un)substituted benzothiazolyl or Ph]. (4-S)-4-

methyl-2-methylthio-4-phenyl-1-phenylamino--2-imidazolin-5-one is representative of I. N1-[(R)-1-(6-fluoro-2-benzothiazolyl)ethyl]-N2- isopropoxycarbonyl-L-valinamide and iso-Pr [2-methyl-1-(phenylethylcarbamoyl)propyl]carbamate are representative of II.

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

L37 ANSWER 2 OF 8 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1998:682071 CAPLUS Full-text

DOCUMENT NUMBER: 129:299238

ORIGINAL REFERENCE NO.: 129:60941a,60944a

TITLE: Synergistic fungicidal compositions containing a

INVENTOR(S): 3-phenylpyrazole derivative  
Chazalet, Maurice; Gouet, Jean-Marie;  
Peignier, Raymond

PATENT ASSIGNEE(S): Rhone-Poulenc Agro, Fr.

SOURCE: PCT Int. Appl., 62 pp.

CODEN: PIXXD2


DOCUMENT TYPE: Patent

LANGUAGE: French

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9843480	A1	19981008	WO 1998-FR608	
19980326 <--				
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
AU 9870512	A	19981022	AU 1998-70512	
19980326 <--				
PRIORITY APPLN. INFO.:			FR 1997-4101	A
19970328 <--			WO 1998-FR608	W
19980326 <--				
OTHER SOURCE(S):	MARPAT	129:299238		
GI				



AB The invention concerns fungicide comps. containing a 3-phenylpyrazole I (X1-5 = H, halo, nitro or alkyl; two of the adjacent X1-5 can further form with the Ph to which they are bound 2,2-difluorobenzodioxolyl; provided that X1-5 cannot each be H at the same time) mixed with a known fungicide. 4-Chloro-3-(3,5-dichlorophenyl)-1H-pyrazole is the preferred I.

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L37 ANSWER 3 OF 8 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 1996:425533 CAPLUS Full-text  
 DOCUMENT NUMBER: 125:79394  
 ORIGINAL REFERENCE NO.: 125:14931a,14934a  
 TITLE: Lawn fungicide  
 INVENTOR(S): Chazalet, Maurice; Gouet, Jean Marie; White, Mark  
 PATENT ASSIGNEE(S): Rhone Poulenc Agrochimie, Fr.  
 SOURCE: Fr. Demande, 8 pp.  
 CODEN: FRXXBL  
 DOCUMENT TYPE: Patent  
 LANGUAGE: French  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 2726737	A1	19960515	FR 1995-12891	
19951026 <--				
FR 2726737	B1	19970704		
PRIORITY APPLN. INFO.:			FR 1995-12891	
19951026 <--				

AB Triticonazole is a lawn fungicide. especially active against Sclerotinia, Puccinia Laetisaria, Fusarium and Gaeumannomyces on Poa, Agrostis, Festuca, Phleum, Lolium and Zoysia.

REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L37 ANSWER 4 OF 8 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 1995:687041 CAPLUS Full-text  
 DOCUMENT NUMBER: 123:77184  
 ORIGINAL REFERENCE NO.: 123:13587a,13590a

TITLE: Synergistic combinations of a fungicide having  
anazole group with an insecticide having a  
pyrazole,  
pyrrole or phenylimidazole group.  
INVENTOR(S): Colliot, Francois; Guot, Jean-Marie; Molle,  
Francis; Duvert, Patrice  
PATENT ASSIGNEE(S): Rhone Poulenc Agrochimie, Fr.  
SOURCE: PCT Int. Appl., 29 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: French  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9512314	A1	19950511	WO 1994-FR1254	
19941027 <--				
W: AU, BR, BY, CA, CN, JP, KR, KZ, PL, RO, RU, SI, UA, US, VN				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL,				
PT, SE				
FR 2711893	A1	19950512	FR 1993-13400	
19931104 <--				
FR 2711893	B1	19960112		
FR 2712144	A1	19950519	FR 1994-11214	
19940914 <--				
FR 2712144	B1	19970718		
CA 2175818	A1	19950511	CA 1994-2175818	
19941027 <--				
AU 9481094	A	19950523	AU 1994-81094	
19941027 <--				
AU 690160	B2	19980423		
EP 726709	A1	19960821	EP 1995-900169	
19941027 <--				
EP 726709	B1	19971203		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL,				
PT, SE				
CN 1140976	A	19970122	CN 1994-194753	
19941027 <--				
CN 1078043	C	20020123		
JP 09504538	T	19970506	JP 1995-513044	
19941027 <--				
BR 9408163	A	19971028	BR 1994-8163	
19941027 <--				
AT 160672	T	19971215	AT 1995-900169	
19941027 <--				
ES 2110308	T3	19980201	ES 1995-900169	
19941027 <--				
RU 2141203	C1	19991120	RU 1996-112105	
19941027 <--				
RO 115930	B1	20000830	RO 1996-928	
19941027 <--				
PL 180374	B1	20010131	PL 1994-314183	
19941027 <--				
ZA 9408725	A	19950703	ZA 1994-8725	

19941104 <--  
 CN 1108043 A 19950913 CN 1994-117809  
 19941104 <--  
 US 5877194 A 19990302 US 1997-953318  
 19971017 <--  
 PRIORITY APPLN. INFO.: FR 1993-13400 A  
 19931104 <--  
 FR 1994-11214 A  
 19940914 <--  
 WO 1994-FR1254 W  
 19941027 <--  
 US 1996-640828 B1  
 19960801 <--  
 AB Agrochem. combinations contain a fungicide having an azole group,  
 such as triticonazole, and an insecticide having a pyrazole,  
 pyrrole or phenylimidazole group, such as fipronil. The method  
 may include applying a single composition containing both active  
 substances or applying two compns. each containing one of the  
 active substances, either at the same time, or one after the  
 other, to achieve a combined effect.  
 REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE  
 FOR THIS  
 REFORMAT RECORD. ALL CITATIONS AVAILABLE IN THE

L37 ANSWER 5 OF 8 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 1995:267266 CAPLUS Full-text  
 DOCUMENT NUMBER: 122:25878  
 ORIGINAL REFERENCE NO.: 122:5021a,5024a  
 TITLE: Improving the vigor and health of plants, such  
 as  
 cereals, with triazole derivatives.  
 INVENTOR(S): Gatineau, Francis; Gout, Jean-Marie;  
 Leroux, Bernard  
 PATENT ASSIGNEE(S): Rhone-Poulenc Agrochimie, Fr.  
 SOURCE: Eur. Pat. Appl.  
 CODEN: EPXXDW  
 DOCUMENT TYPE: Patent  
 LANGUAGE: French  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 622020	A1	19941102	EP 1994-420127	
19940422 <--				
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE				
FR 2704388	A1	19941104	FR 1993-5193	
19930427 <--				
FR 2704388	B1	19950609		
ZA 9402896	A	19950104	ZA 1994-2896	
19940426 <--				
HU 71060	A2	19951128	HU 1994-1189	
19940426 <--				
CA 2122331	A1	19941028	CA 1994-2122331	

19940427 <--  
 PRIORITY APPLN. INFO.: FR 1993-5193 A  
 19930427 <--  
 OTHER SOURCE(S): MARPAT 122:25878  
 AB Seed treatment with a triazole derivative (Markush given),  
 specifically triticonazole, improves the vigor and health of  
 cereals. The seeds are optionally post-treated with cycocel or  
 Ethephon.

L37 ANSWER 6 OF 8 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 1988:112454 CAPLUS Full-text  
 DOCUMENT NUMBER: 108:112454  
 ORIGINAL REFERENCE NO.: 108:18425a,18428a  
 TITLE: Preparation of  
 4-bromo-2-cyano-6-(trifluoromethyl)-1H-  
 benzimidazole-1-  
 sulfonamides as agrochemical fungicides  
 INVENTOR(S): Souche, Jean Luc; Gouot, Jean Marie  
 PATENT ASSIGNEE(S): Rhone-Poulenc Agrochimie, Fr.  
 SOURCE: Fr. Demande, 23 pp.  
 CODEN: FRXXBL  
 DOCUMENT TYPE: Patent  
 LANGUAGE: French  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	----
FR 2594437	A1	19870821	FR 1986-2455	
19860219 <--				
DD 260211	A5	19880921	DD 1987-299924	
19870216 <--				
AU 8768860	A	19870820	AU 1987-68860	
19870217 <--				
EP 239508	A2	19870930	EP 1987-420046	
19870217 <--				
EP 239508	A3	19871014		
R: AT, BE, CH, DE, ES, FR, GB, GR, IT, LI, LU, NL, SE				
ZA 8701141	A	19870930	ZA 1987-1141	
19870217 <--				
DK 8700812	A	19870820	DK 1987-812	
19870218 <--				
FI 8700669	A	19870820	FI 1987-669	
19870218 <--				
NO 8700639	A	19870820	NO 1987-639	
19870218 <--				
JP 62205063	A	19870909	JP 1987-35441	
19870218 <--				
HU 43318	A2	19871028	HU 1987-629	
19870218 <--				
BR 8700779	A	19871222	BR 1987-779	
19870219 <--				
PRIORITY APPLN. INFO.:			FR 1986-2455	A
19860219 <--				
OTHER SOURCE(S):		CASREACT 108:112454		

GI



AB The title compds. (I; R = C2-4 dialkylamino) were prepared as plant fungicides. 2,4-O2N(F3C)C6H3NH2 was brominated and reduced with SnCl2 to give 3-bromo-5-(trifluoromethyl)-1,2-benzenediamine-HCl which was cyclocondensed with Cl3CCO2Me to give 4-bromo-2-(trichloromethyl)-6-(trifluoromethyl)-1H-benzimidazole. The latter was treated with aqueous NH3 to give the 2-cyano analog which was stirred at 20° with a suspension of K in acetone while Me2NSO2Cl was slowly added to give I (R = Me2N) (II). Potato plants infected with Phytophthora infestans were sprayed at 10 day intervals with a spray containing 15 g II/hL at an application rate of 1000 L/ha. Four days after the 4th application 2.7% of the leaf surface showed fungal attack, compared to 30% using another, known benzimidazolesulfonamide fungicide.

REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

L37 ANSWER 7 OF 8 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 1987:551506 CAPLUS Full-text  
 DOCUMENT NUMBER: 107:151506  
 ORIGINAL REFERENCE NO.: 107:24325a,24328a  
 TITLE: Differential diagnosis of fungal diseases in cereals  
 INVENTOR(S): Gouet, Jean Marie; Paviot, Jean  
 PATENT ASSIGNEE(S): Rhone-Poulenc Agrochimie, Fr.  
 SOURCE: Belg., 15 pp.  
 CODEN: BEXXAL  
 DOCUMENT TYPE: Patent  
 LANGUAGE: French  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
BE 905482	A1	19870324	BE 1986-217203	
19860924 <--				
DK 8604547	A	19870326	DK 1986-4547	
19860924 <--				
GB 2180853	A	19870408	GB 1986-23047	
19860925 <--				

GB 2180853 B 19891213 A  
PRIORITY APPLN. INFO.: FR 1985-14403 A  
19850925 <--

AB A method for the differential diagnosis of *Pseudocercospora* herpotrichoides, *Rhizoctonia cerealis* and *Fusarium* consists in contacting cereal stem segments with 3 in vitro culture media, each containing a fungal growth inhibitor specific for the pertinent species. Three petri dishes were filled with the PDA medium containing 100 ppm streptomycin, 50 ppm penicillin, and 50 ppm aureomycin. The 1st dish, for the differential diagnosis of *P. herpotrichoides*, contained 200 ppm ditalimphos and 5 ppm iprodione. The 2nd dish, for *R. cerealis*, contained 0.5 ppm carbendazim and 0.5 ppm prochloras. The 3rd dish, for *Fusarium*, contained 2 ppm penconazol and 5 ppm iprodione.

L37 ANSWER 8 OF 8 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1981:509820 CAPLUS Full-text  
DOCUMENT NUMBER: 95:109820  
ORIGINAL REFERENCE NO.: 95:18345a, 18348a  
TITLE: Pentachloronitrobenzene metabolism in peanut.  
3.  
Metabolism in peanut cell suspension cultures  
AUTHOR(S): Lamoureux, Gerald L.; Gouot, Jean Marie;  
Davis, David G.; Rusness, Donald G.  
CORPORATE SOURCE: Metab. Radiat. Res. Lab., Sci. Educ. Adm.,  
Fargo, ND, 58105, USA  
SOURCE: Journal of Agricultural and Food Chemistry ( 1981), 29(5), 996-1002  
CODEN: JAFCAU; ISSN: 0021-8561  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
GI



AB The metabolism of U-14C-labeled PCNB (I) [82-68-8] was studied in peanut (*Arachis hypogaea*) cell suspension cultures over a 14-day period. The primary metabolic pathways involved an initial conjugation with glutathione. Seven major metabolites were detected by high-performance liquid chromatog., and 5 of these were identified by mass spectrometry of suitable derivs.: S-(pentachlorophenyl)glutathione [75005-81-1], S-(ar-tetrachloronitrophenyl)-N-malonylcysteine [74998-44-0], and S-(pentachlorophenyl)-N-malonylcysteine [75005-77-5]. Several precursor-product relationships were demonstrated. Nonextractable residue, S-(pentachlorophenyl)-N-malonylcysteine, S-(ar-



tetrachloronitrophenyl)-N-malonylcysteine, and metabolite III appeared to be terminal metabolic products. PCNB metabolism in peanut cell suspension cultures was compared to PCNB metabolism in the roots of intact peanut plants. The primary differences between the 2 systems appeared to be quant. Pentachloroaniline [527-20-8] and nonextractable residue were produced in larger amts. in intact peanut plants than in the cell suspension cultures. Several advantages and disadvantages of conducting metabolism studies in cell suspension cultures were discussed.

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate substance identification.

```
=> s ?benzamide? and (respirat? or electrons and transport)
    35699 ?BENZAMIDE?
    224919 RESPIRAT?
    283950 ELECTRONS
    818418 TRANSPORT
    7155 TRANSPORTS
    821378 TRANSPORT
        (TRANSPORT OR TRANSPORTS)
L1      565 ?BENZAMIDE? AND (RESPIRAT? OR ELECTRONS AND TRANSPORT)

=> s l1 and (py<2003 or ay<2003 or pry<2003)
    22983367 PY<2003
    4503987 AY<2003
    3972879 PRY<2003
L2      278 L1 AND (PY<2003 OR AY<2003 OR PRY<2003)

=> s l2 and (?fung? or ?herb? or ?pest?)
    263920 ?FUNG?
    160492 ?HERB?
    150744 ?PEST?
L3      23 L2 AND (?FUNG? OR ?HERB? OR ?PEST?)

=> d l3 ibib abs ti hit 1-10

L3  ANSWER 1 OF 23  CAPLUS  COPYRIGHT 2009 ACS on STN
ACCESSION NUMBER:  2008:1280175  CAPLUS  Full-text
DOCUMENT NUMBER:  149:506150
TITLE:
    Phenoxypropionic acid benzamides and related
modulators        compounds as selective androgen receptor
                    (SARMs) for treating diabetes, diseases
associated
                    with diabetes, and other disorders
INVENTOR(S):      Dalton, James T.; Miller, Duane D.
PATENT ASSIGNEE(S):  University of Tennessee Research Foundation,
USA
SOURCE:           PCT Int. Appl., 194pp.
                   CODEN: PIXXD2
DOCUMENT TYPE:     Patent
LANGUAGE:          English
FAMILY ACC. NUM. COUNT:  40
PATENT INFORMATION:
```

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2008127717	A1	20081023	WO 2008-US4816	
20080414				
W: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW				
HR, HU, SI, SK, SN, TD, ZM, ZW,				
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
EP 1487780	A1	20041222	EP 2003-713345	
20030224				
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
AU 2006201538	A1	20060504	AU 2006-201538	
20060412 <--				
AU 2006201538	B2	20080221		
US 20070281906	A1	20071206	US 2007-785064	
20070413 <--				
AU 2007201969	A1	20070524	AU 2007-201969	
20070503				
AU 2007202324	A1	20070614	AU 2007-202324	
20070522 <--				
AU 2008201811	A1	20080515	AU 2008-201811	
20080424				
AU 2008202236	A1	20080612	AU 2008-202236	
20080521				
PRIORITY APPLN. INFO.:			US 2007-785064	A
20070413				
19990607 <--			AU 2002-2002364949	A3
20010823 <--			AU 2001-285230	A3
20010823 <--			AU 2001-85230	T0
20011206 <--			US 2001-336185P	P
			US 2002-310150	A2

20021205 <--

20030224

20030224

20031014

20031014

20031216

20040607

20041012

20041216

20050607

20050831

20050907

20060216

20060412

20060817

20060817

20060828

20061206

OTHER SOURCE(S):  
GI

MARPAT 149:506150

AU 2003-216153

A3

WO 2003-US3122

W

AU 2003-287074

A3

US 2003-510138P

P

US 2003-529573P

P

US 2004-861923

A2

US 2004-961380

A2

US 2004-13214

A2

US 2005-146427

A2

US 2005-712390P

P

US 2005-220414

A2

US 2006-355187

A2

AU 2006-201538

A3

US 2006-505363

A2

US 2006-505499



A2

US 2006-510844

A2

US 2006-634380

A2



AB This invention provides use of a SARM compound or a composition comprising the same in treating a variety of diseases or conditions in a subject, including, inter-alia, a diabetes

disease, and/or disorder such as cardiovascular disease, atherosclerosis, cerebrovascular conditions, diabetic nephropathy, diabetic neuropathy, and diabetic retinopathy. The SARMs have general formula I (wherein X = bond, O, CH<sub>2</sub>, NH, etc.; T = OH, OR, NHAc, NHCOR; Z = NO<sub>2</sub>, cyano, CO<sub>2</sub>H, COR, CONHR; Y = H, alkoxy, CF<sub>3</sub>, etc.; Q = alkyl, halo, cyano, etc.; R = alkyl, haloalkyl, etc.; R<sub>1</sub> = Me, CF<sub>3</sub>, etc.). II is the compound of prime interest in the patent. I can be formulated alone or with other drugs.

TI Phenoxypropionic acid benzamides and related compounds as selective androgen receptor modulators (SARMs) for treating diabetes, diseases associated with diabetes, and other disorders  
REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

TI Phenoxypropionic acid benzamides and related compounds as selective androgen receptor modulators (SARMs) for treating diabetes,

diseases associated with diabetes, and other disorders

PRAI US 2007-785064 A 20070413

AU 2002-2002364949 A3 19990607 <--

AU 2001-285230 A3 20010823 <--

AU 2001-85230 T0 20010823 <--

US 2001-336185P P 20011206 <--

US 2002-310150 A2 20021205 <--

AU 2003-216153 A3 20030224

WO 2003-US3122 W 20030224

AU 2003-287074 A3 20031014

US 2003-510138P P 20031014

US 2003-529573P P 20031216

US 2004-861923 A2 20040607

US 2004-961380 A2 20041012

US 2004-13214 A2 20041216

US 2005-146427 A2 20050607

US 2005-712390P P 20050831

US 2005-220414 A2 20050907

US 2006-355187 A2 20060216

AU 2006-201538 A3 20060412

US 2006-505363 A2 20060817

US 2006-505499 A2 20060817

US 2006-510844 A2 20060828

US 2006-634380 A2 20061206

L3 ANSWER 2 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2007:1396630 CAPLUS Full-text

DOCUMENT NUMBER: 148:45855

TITLE: Phenoxypropionic acid benzamides and related compounds as selective androgen receptor

modulators

(SARMs) for treating diabetes, diseases

associated

with diabetes, and other disorders

INVENTOR(S): Dalton, James T.; Miller, Duane D.

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 102 pp., Cont.-in-part of U.S.

Ser. No. 634,380.

CODEN: USXXCO

DOCUMENT TYPE:

Patent

LANGUAGE:

English

FAMILY ACC. NUM. COUNT:

40

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	----
20070413 <--	A1	20071206	US 2007-785064	
20021205 <--	A1	20040506	US 2002-310150	
20030224	A1	20041222	EP 2003-713345	
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
20041012	A1	20060126	US 2004-961380	
20041216	A1	20060112	US 2004-13214	
20050607 <--	A1	20060216	US 2005-146427	
20050907 <--	A1	20060525	US 2005-220414	
20060216 <--	A1	20061012	US 2006-355187	
20060412 <--	A1	20060504	AU 2006-201538	
20060817 <--	B2	20080221		
20060828	A1	20070531	US 2006-505499	
20070503	A1	20070726	US 2006-505363	
20070522 <--	A1	20070322	US 2006-510844	
20080414	A1	20070712	US 2006-634380	
20080414	A1	20070524	AU 2007-201969	
20080414	A1	20070614	AU 2007-202324	
20080414	A1	20081023	WO 2008-US4816	
W: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,				
CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES,				
FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE,				
KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,				
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH,				
PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY,				

TJ, TM,  
 TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW  
 RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR,  
 HR, HU,  
 IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE,  
 SI, SK,  
 TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE,  
 SN, TD,  
 TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG,  
 ZM, ZW,  
 AM, AZ, BY, KG, KZ, MD, RU, TJ, TM  
 AU 2008201811 A1 20080515 AU 2008-201811  
 20080424  
 AU 2008202236 A1 20080612 AU 2008-202236  
 20080521  
 PRIORITY APPLN. INFO.: US 2001-336185P P  
 20011206 <-- US 2002-310150 A2  
 20021205 <-- US 2003-510138P P  
 20031014 US 2003-529573P P  
 20031216 US 2004-861923 A2  
 20040607 US 2004-961380 A2  
 20041012 US 2004-13214 A2  
 20041216 US 2005-146427 A2  
 20050607 US 2005-712390P P  
 20050831 US 2005-220414 A2  
 20050907 US 2006-355187 A2  
 20060216 US 2006-505363 A2  
 20060817 US 2006-505499 A2  
 20060817 US 2006-510844 A2  
 20060828 US 2006-634380 A2  
 20061206 AU 2002-2002364949 A3  
 19990607 <-- US 2000-367355P P  
 20000824 <-- US 2001-300083P P  
 20010625 <-- AU 2001-285230 A3  
 20010823 <-- AU 2001-85230 T0  
 20010823 <-- US 2001-935044 A2  
 20010823 <--

20010823 <--	US 2001-935045	A2
20021015 <--	US 2002-270232	A2
20021015 <--	US 2002-418166P	P
20021022 <--	US 2002-277108	A2
20030224	AU 2003-216153	A3
20030224	US 2003-371213	A2
20030224	WO 2003-US3122	W
20031014	AU 2003-287074	A3
20031014	US 2003-683157	A2
20040609	US 2004-863524	A2
20040928	US 2004-613206P	P
20050223	US 2005-62752	A2
20050510	US 2005-125159	A2
20060214	US 2006-353225	A2
20060412	AU 2006-201538	A3
20070413	US 2007-785064	A

AB This invention provides use of a SARM compound or a composition comprising the same in treating a variety of diseases or conditions in a subject, including, inter-alia, a diabetes disease and/or disorder such as cardiovascular disease, atherosclerosis, cerebrovascular conditions, diabetic nephropathy, diabetic neuropathy and diabetic retinopathy. The SARMS have general formula I (wherein X = bond, O, CH<sub>2</sub>, NH, etc.; T = OH, OR, NHAc, NHCOR; Z = NO<sub>2</sub>, cyano, CO<sub>2</sub>H, COR, CONHR; Y = H, alkoxy, CF<sub>3</sub>, etc.; Q = alkyl, halo, cyano, etc.; R = alkyl, haloalkyl, etc.; R<sub>1</sub> = Me, CF<sub>3</sub>, etc.). II is the compound of prime interest in the patent. I can be formulated alone or with other drugs.

TI Phenoxypropionic acid benzamides and related compounds as selective androgen receptor modulators (SARMS) for treating diabetes,

diseases associated with diabetes, and other disorders

TI Phenoxypropionic acid benzamides and related compounds as selective androgen receptor modulators (SARMS) for treating diabetes,

diseases associated with diabetes, and other disorders

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	----
PI US 20070281906	A1	20071206	US 2007-785064	

20070413 <--				
US 20040087557	A1	20040506	US	2002-310150
20021205 <--				
EP 1487780	A1	20041222	EP	2003-713345
20030224				
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
US 20060019931	A1	20060126	US	2004-961380
20041012				
US 20060009488	A1	20060112	US	2004-13214
20041216				
US 20060035965	A1	20060216	US	2005-146427
20050607 <--				
US 20060111441	A1	20060525	US	2005-220414
20050907 <--				
US 20060229362	A1	20061012	US	2006-355187
20060216 <--				
AU 2006201538	A1	20060504	AU	2006-201538
20060412 <--				
AU 2006201538	B2	20080221		
US 20070123563	A1	20070531	US	2006-505499
20060817 <--				
US 20070173546	A1	20070726	US	2006-505363
20060817 <--				
US 20070066568	A1	20070322	US	2006-510844
20060828				
US 20070161608	A1	20070712	US	2006-634380
20061206 <--				
AU 2007201969	A1	20070524	AU	2007-201969
20070503				
AU 2007202324	A1	20070614	AU	2007-202324
20070522 <--				
WO 2008127717	A1	20081023	WO	2008-US4816
20080414				
W: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,				
CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES,				
FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE,				
KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,				
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH,				
PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM,				
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW				
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU,				
IE, IS, IT, LI, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK,				
TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,				
TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW,				
AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				



20080424	AU 2008201811	A1	20080515	AU 2008-201811
20080521	AU 2008202236	A1	20080612	AU 2008-202236
PRAI	US 2001-336185P	P	20011206	<--
	US 2002-310150	A2	20021205	<--
	US 2003-510138P	P	20031014	
	US 2003-529573P	P	20031216	
	US 2004-861923	A2	20040607	
	US 2004-961380	A2	20041012	
	US 2004-13214	A2	20041216	
	US 2005-146427	A2	20050607	
	US 2005-712390P	P	20050831	
	US 2005-220414	A2	20050907	
	US 2006-355187	A2	20060216	
	US 2006-505363	A2	20060817	
	US 2006-505499	A2	20060817	
	US 2006-510844	A2	20060828	
	US 2006-634380	A2	20061206	
	AU 2002-2002364949	A3	19990607	<--
	US 2000-367355P	P	20000824	<--
	US 2001-300083P	P	20010625	<--
	AU 2001-285230	A3	20010823	<--
	AU 2001-85230	T0	20010823	<--
	US 2001-935044	A2	20010823	<--
	US 2001-935045	A2	20010823	<--
	US 2002-270232	A2	20021015	<--
	US 2002-418166P	P	20021015	<--
	US 2002-277108	A2	20021022	<--
	AU 2003-216153	A3	20030224	
	US 2003-371213	A2	20030224	
	WO 2003-US3122	W	20030224	
	AU 2003-287074	A3	20031014	
	US 2003-683157	A2	20031014	
	US 2004-863524	A2	20040609	
	US 2004-613206P	P	20040928	
	US 2005-62752	A2	20050223	
	US 2005-125159	A2	20050510	
	US 2006-353225	A2	20060214	
	AU 2006-201538	A3	20060412	
	US 2007-785064	A	20070413	

L3 ANSWER 3 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 2006:1256641 CAPLUS Full-text  
 DOCUMENT NUMBER: 146:50262  
 TITLE: Antibiotic kit and compositions  
 INVENTOR(S): Friedman, Doron; Besonov, Alex; Tamarkin, Dov;  
 Eini, Meir  
 PATENT ASSIGNEE(S): Foamix Ltd., Israel  
 SOURCE: U.S. Pat. Appl. Publ., 31pp., Cont.-in-part of  
 U.S. Ser. No. 532,618.  
 CODEN: USXXCO  
 DOCUMENT TYPE: Patent

LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 33  
 PATENT INFORMATION:

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	-----	----	-----	-----	-----
20060607 <--	US 20060269485	A1	20061130	US 2006-448490	
20031024 <--	WO 2004037225	A2	20040506	WO 2003-IB5527	
	WO 2004037225	A3	20041229		
CH, CN,	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,				
GE, GH,	CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD,				
LK, LR,	GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,				
OM, PH,	LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ,				
TT, TZ,	PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR,				
	UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
AZ, BY,	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM,				
EE, ES,	KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,				
SK, TR,	FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI,				
TD, TG	BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN,				
20040804	US 20050069566	A1	20050331	US 2004-911367	
20051222 <--	US 20060140984	A1	20060629	US 2005-532618	
20060607	AU 2006339311	A2	20070907	AU 2006-339311	
	AU 2006339311	A1	20070907		
	CA 2611577	A1	20070907	CA 2006-2611577	
20060607	WO 2007099396	A2	20070907	WO 2006-IB3975	
20060607	WO 2007099396	A3	20080313		
CA, CH,	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,				
GB, GD,	CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,				
KP, KR,	GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN,				
MW, MX,	KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN,				
SC, SD,	MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU,				
US, UZ,	SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG,				
	VC, VN, ZA, ZM, ZW				
	RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR,				

HU, IE,  
 BF, BJ,  
 BW, GH,  
 AZ, BY,  
 IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR,  
 CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG,  
 GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM,  
 KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA  
 EP 1919449 A2 20080514 EP 2006-847249  
 20060607  
 R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR,  
 HU, IE,  
 IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK,  
 TR, AL,  
 BA, HR, MK, RS  
 US 20070292355 A1 20071220 US 2007-732547  
 20070404 <--  
 IN 2007KN04925 A 20080704 IN 2007-KN4925  
 20071218  
 PRIORITY APPLN. INFO.: US 2002-429546P P  
 20021129 <--  
 US 2003-492385P P  
 20030804  
 WO 2003-IB5527 W  
 20031024  
 US 2004-911367 A2  
 20040804  
 US 2005-688244P P  
 20050607  
 US 2005-532618 A2  
 20051222  
 IL 2002-152486 A  
 20021025 <--  
 US 2003-497648P P  
 20030825  
 US 2003-530015P P  
 20031216  
 US 2004-835505 A2  
 20040428  
 US 2004-922358 A2  
 20040820  
 US 2005-41921 A2  
 20050124  
 US 2006-789186P P  
 20060404  
 US 2006-448490 A2  
 20060607  
 WO 2006-IB3975 W  
 20060607  
 US 2006-861620P P  
 20061129  
 US 2007-880434P P  
 20070112  
 AB The present invention relates to a therapeutic kit to provide an  
 effective dosage of an antibiotic including an aerosol packaging  
 assembly. The assembly includes a container accommodating a  
 pressurized product; and an outlet capable of releasing the



pressurized product as a foam, wherein the pressurized product comprises a foamable composition of an antibiotic; at least one organic carrier selected from the group consisting of a hydrophobic organic carrier, an organic polar solvent, an emollient and mixts. at 2-50%, a surfactant, 0.01-5% by weight of at least one polymeric additive selected from the group consisting of a bioadhesive agent, a gelling agent, a film forming agent and a phase change agent, water; and liquefied or compressed gas propellant at 3-25% by weight of the total composition

TI Antibiotic kit and compositions  
 PRAI US 2002-429546P P 20021129 <--  
 US 2003-492385P P 20030804  
 WO 2003-IB5527 W 20031024  
 US 2004-911367 A2 20040804  
 US 2005-688244P P 20050607  
 US 2005-532618 A2 20051222  
 IL 2002-152486 A 20021025 <--  
 US 2003-497648P P 20030825  
 US 2003-530015P P 20031216  
 US 2004-835505 A2 20040428  
 US 2004-922358 A2 20040820  
 US 2005-41921 A2 20050124  
 US 2006-789186P P 20060404  
 US 2006-448490 A2 20060607  
 WO 2006-IB3975 W 20060607  
 US 2006-861620P P 20061129  
 US 2007-880434P P 20070112

L3 ANSWER 4 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 2004:291088 CAPLUS Full-text  
 DOCUMENT NUMBER: 140:321350  
 TITLE: Preparation of indazolecarboxamides as CDK1,  
 CDK2, and CDK4 inhibitors for treating CDK-related  
 diseases, in particular cancer  
 INVENTOR(S): D'Orchymont, Hugues; Van Hijfte, Luc;  
 Zimmermann, Andre  
 PATENT ASSIGNEE(S): Sanofi-Synthelabo, Fr.  
 SOURCE: Fr. Demande, 90 pp.  
 CODEN: FRXXBL  
 DOCUMENT TYPE: Patent  
 LANGUAGE: French  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	----
FR 2845382	A1	20040409	FR 2002-12188	
20021002 <--				
WO 2004031158	A1	20040415	WO 2003-FR2862	
20030930 <--				
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,				

GD, GE, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB,  
 LC, LK, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ,  
 NO, NZ, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI,  
 TJ, TM, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY,  
 TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW  
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM,  
 AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,  
 EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI,  
 SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN,  
 TD, TG AU 2003299125 A1 20040423 AU 2003-299125  
 20030930 <--  
 EP 1549620 A1 20050706 EP 2003-798949  
 20030930 <--  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,  
 MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK  
 JP 2006504711 T 20060209 JP 2004-540862  
 20030930 <--  
 US 20060004000 A1 20060105 US 2005-96375  
 20050401 <--  
 US 7482342 B2 20090127  
 PRIORITY APPLN. INFO.: FR 2002-12188 A  
 20021002 <-- WO 2003-FR2862 W  
 20030930  
 OTHER SOURCE(S): MARPAT 140:321350  
 GI



AB Title compds. I [R1 = H, halo, NH2, NHR2, NHCOR2, NO2, CN, CH2NH2, CH2NHR2, (un)substituted Ph, heteroaryl; Ar = (un)substituted Ph, heteroaryl; R2 = Ph, heteroaryl, (un)substituted alkyl (substituent = Ph or heteroaryl); n = 0, 1, 2, or 3; PG = protecting group selected from trimethylsilylethoxymethyl, mesitylenesulfonyl; their free bases, addition salts with acids, solvates and hydrates; with the exclusion of certain compds.] were prepared as cyclin-dependent kinase (CDK)-1, CDK2, and CDK4 inhibitors for treating cdk-related diseases, in particular cancer. For instance, reacting indazole-3-carboxylic acid with N-phenyl-1,4-phenylenediamine in the presence of DCC gave 58% II. I displayed IC50 values < 20 µM for the inhibition of CDK2, CDK1, and CDK4 in a test for measuring the enzymic activity of CDK2/Cyclin A, CDK1/Cyclin B, and CDK4/Cyclin D1, resp. I are useful for treating cancers, autoimmune diseases, inflammations, cardiovascular diseases, viral and fungal infections, hematol. diseases, and degenerative diseases of muscular system.

TI Preparation of indazolecarboxamides as CDK1, CDK2, and CDK4 inhibitors for

treating CDK-related diseases, in particular cancer

REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

# RE FORMAT

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 2845382	A1	20040409	FR 2002-12188	
20021002 <--				
WO 2004031158	A1	20040415	WO 2003-FR2862	
20030930 <--				
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
AU 2003299125	A1	20040423	AU 2003-299125	
20030930 <--				
EP 1549620	A1	20050706	EP 2003-798949	
20030930 <--				
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,			

IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK  
 JP 2006504711 T 20060209 JP 2004-540862  
 20030930 <--  
 US 20060004000 A1 20060105 US 2005-96375  
 20050401 <--  
 US 7482342 B2 20090127  
 PRAI FR 2002-12188 A 20021002 <--  
 WO 2003-FR2862 W 20030930

AB Title compds. I [R1 = H, halo, NH2, NHR2, NHCOR2, NO2, CN, CH2NH2, CH2NHR2, (un)substituted Ph, heteroaryl; Ar = (un)substituted Ph, heteroaryl; R2 = Ph, heteroaryl, (un)substituted alkyl (substituent = Ph or heteroaryl); n = 0, 1, 2, or 3; PG = protecting group selected from trimethylsilylethoxymethyl, mesitylenesulfonyl; their free bases, addition salts with acids, solvates and hydrates; with the exclusion of certain compds.] were prepared as cyclin-dependent kinase (CDK)-1, CDK2, and CDK4 inhibitors for treating cdk-related diseases, in particular cancer. For instance, reacting indazole-3-carboxylic acid with N-phenyl-1,4-phenylenediamine in the presence of DCC gave 58% II. I displayed IC50 values < 20 µM for the inhibition of CDK2, CDK1, and CDK4 in a test for measuring the enzymic activity of CDK2/Cyclin A, CDK1/Cyclin B, and CDK4/Cyclin D1, resp. I are useful for treating cancers, autoimmune diseases, inflammations, cardiovascular diseases, viral and fungal infections, hematol. diseases, and degenerative diseases of muscular system


L3 ANSWER 5 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2004:266876 CAPLUS Full-text  
 DOCUMENT NUMBER: 140:287180  
 TITLE: Preparation of arylamines, arylamides and arylureas as

inhibitors of undesired cell proliferation  
 Knolle, Jochen; Schutkowski, Mike; Hummel, Gerd  
 INVENTOR(S):  
 Gerd  
 PATENT ASSIGNEE(S): Jerini Ag, Germany  
 SOURCE: Eur. Pat. Appl., 126 pp.  
 CODEN: EPXXDW  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1402887	A1	20040331	EP 2002-20922	
20020918 <--				
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
WO 2004030664	A2	20040415	WO 2003-EP10415	
20030918 <--				
WO 2004030664	A3	20040812		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,				
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,				

GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
 LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO,  
 NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ,  
 TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW  
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM,  
 AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,  
 EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI,  
 SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN,  
 TD, TG  
 AU 2003277871 A1 20040423 AU 2003-277871  
 20030918 <--  
 PRIORITY APPLN. INFO.: EP 2002-20922 A  
 20020918 <--  
 WO 2003-EP10415 W  
 20030918  
 OTHER SOURCE(S): MARPAT 140:287180  
 GI



II

AB Title compds. A-X-Y [A = cycloalkyl, heterocyclyl, aryl, etc.; X = [(CRaRb)nNRcCONR'(CRaRb)m]p, etc; n, m = 0-10 provided that if n = 0, m is not 0; p = 0-10; Ra-c, R' = H, alkyl, cycloalkyl, etc.; Y = alkyl, cycloalkyl, etc.; I] are prepared For instance, 6-amino-2,4-dichloro-3-methylphenol•HCl is reacted with 1-adamantylisocyanate (DMSO) to give II. Selected examples of I exhibited cytotoxicity in selected cell lines. I are useful for the treatment of disease that involves abnormal cell proliferation, an undesired cell proliferation, an abnormal mitosis and/or an undesired mitosis.  
 TI Preparation of arylamines, arylamides and arylureas as inhibitors of

undesired cell proliferation  
 REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 1402887	A1	20040331	EP 2002-20922	




20020918 <--  
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,  
MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK  
WO 2004030664 A2 20040415 WO 2003-EP10415  
20030918 <--  
WO 2004030664 A3 20040812  
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,  
CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD,  
GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO,  
NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ,  
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW  
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM,  
AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,  
EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI,  
SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN,  
TD, TG AU 2003277871 A1 20040423 AU 2003-277871  
20030918 <--  
PRAI EP 2002-20922 A 20020918 <--  
WO 2003-EP10415 W 20030918  
IT Respiratory distress syndrome  
(acute; preparation of arylamines, arylamides and arylureas as  
inhibitors of  
undesired cell proliferation)

L3 ANSWER 6 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 2004:80685 CAPLUS Full-text  
DOCUMENT NUMBER: 140:146011  
TITLE: Preparation of bicyclic piperidine derivatives  
as  
antagonists of the CCR1 chemokine receptor  
INVENTOR(S): Blumberg, Laura Cook; Brown, Matthew Frank;  
Hayward,  
Matthew Merrill; Poss, Christopher Stanley  
PATENT ASSIGNEE(S): Pfizer Products Inc., USA  
SOURCE: PCT Int. Appl., 90 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004009588	A1	20040129	WO 2003-IB3155	

20030707 <--  
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,  
CH, CN,  
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD,  
GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO,  
NZ, OM,  
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN,  
TR, TT,  
TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW  
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM,  
AZ, BY,  
KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,  
EE, ES,  
FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI,  
SK, TR,  
BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN,  
TD, TG  
CA 2492110 A1 20040129 CA 2003-2492110  
20030707 <--  
AU 2003281527 A1 20040209 AU 2003-281527  
20030707 <--  
BR 2003012699 A 20050426 BR 2003-12699  
20030707 <--  
EP 1525201 A1 20050427 EP 2003-741007  
20030707 <--  
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,  
MC, PT,  
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK  
CN 1668614 A 20050914 CN 2003-817005  
20030707 <--  
JP 2005533845 T 20051110 JP 2004-522638  
20030707 <--  
US 20040063688 A1 20040401 US 2003-616843  
20030708 <--  
IN 2004DN04155 A 20050401 IN 2004-DN4155  
20041228 <--  
MX 2005000757 A 20050419 MX 2005-757  
20050118 <--  
PRIORITY APPLN. INFO.: US 2002-397263P P  
20020718 <--  
WO 2003-1B3155 W  
20030707  
OTHER SOURCE(S): MARPAT 140:146011  
GI



AB The title compds. [I; a = 1-5; b = 0-4; c = 0-1; Q = alkyl; W = aryl, heteroaryl; Y = O, NH, N(alkyl); Z = O, NH, N(alkyl), N(acetyl); R1 = H, halo, CN, NO2, etc.; R2, R3 = H, alkyl, haloalkyl; R4 = alkylene, (CH2)xO(CH2)y (wherein x, y = 1-2); R5 = H, halo, alkyl, etc.; R6 = H, halo, alkyl, etc.], useful as potent and selective inhibitors of MIP-1 $\alpha$ (CCL3) binding to its receptor CCR1 found on inflammatory and immunomodulatory cells (preferably leukocytes and lymphocytes), were prepared E.g., a multi-step synthesis of (trans)-5-chloro-2-(2-[3-(4-fluorophenoxy)-8-azabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy)benzamide was given. All exemplified compds. I had IC50 of <10  $\mu$ M in the chemotaxis assay. Pharmaceutical composition comprising the compound I is claimed.

II Preparation of bicyclic piperidine derivatives as antagonists of the CCR1

chemokine receptor

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

PRAI US 2002-397263P P 20020718 <--  
WO 2003-1B3155 W 20030707

AB The title compds. [I; a = 1-5; b = 0-4; c = 0-1; Q = alkyl; W = aryl, heteroaryl; Y = O, NH, N(alkyl); Z = O, NH, N(alkyl), N(acetyl); R1 = H, halo, CN, NO2, etc.; R2, R3 = H, alkyl, haloalkyl; R4 = alkylene, (CH2)xO(CH2)y (wherein x, y = 1-2); R5 = H, halo, alkyl, etc.; R6 = H, halo, alkyl, etc.], useful as potent and selective inhibitors of MIP-1 $\alpha$ (CCL3) binding to its receptor CCR1 found on inflammatory and immunomodulatory cells (preferably leukocytes and lymphocytes), were prepared E.g., a multi-step synthesis of (trans)-5-chloro-2-(2-[3-(4-fluorophenoxy)-8-azabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy)benzamide was given. All exemplified compds. I had IC50 of <10  $\mu$ M in the chemotaxis assay. Pharmaceutical composition comprising the compound I is claimed.

L3 ANSWER 7 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2004:80652 CAPLUS Full-text

DOCUMENT NUMBER: 140:146007

TITLE: Preparation of piperidinylketones as as selective

inhibitors of macrophage inflammatory protein

1 $\alpha$

(MIP-1 $\alpha$ ) binding to CCR1 chemokine receptors.

INVENTOR(S): Blumberg, Laura Cook; Brown, Matthew Frank; Hayward,

Matthew Merrill; Poss, Christopher Stanley

PATENT ASSIGNEE(S): Pfizer Products Inc., USA

SOURCE: PCT Int. Appl., 62 pp.

CODEN: PIXXD2


DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

## PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
20030707 <--	A1	20040129	WO 2003-IB2876	
WO 2004009550				
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CA 2492651	A1	20040129	CA 2003-2492651	
20030707 <--				
AU 2003242941	A1	20040209	AU 2003-242941	
20030707 <--				
EP 1534677	A1	20050601	EP 2003-765230	
20030707 <--				
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
BR 2003012946	A	20050712	BR 2003-12946	
20030707 <--				
CN 1668592	A	20050914	CN 2003-817092	
20030707 <--				
JP 2005537279	T	20051208	JP 2004-522601	
20030707 <--				
US 20040063759	A1	20040401	US 2003-616844	
20030708 <--				
IN 2004DN04166	A	20070511	IN 2004-DN4166	
20041229 <--				
ZA 2005000067	A	20051102	ZA 2005-67	
20050104 <--				
MX 2005000380	A	20050331	MX 2005-380	
20050106 <--				
PRIORITY APPLN. INFO.:			US 2002-397108P	P
20020718 <--			WO 2003-IB2876	W
20030707				
OTHER SOURCE(S):	MARPAT 140:146007			
GI				



I

AB Title compds. [I; m = 1-5; n = 0-4; p = 0-1; Q = alkyl; W = aryl, heteroaryl; Y = O, NR8; R8 = H, alkyl; Z = O, NR9; R9 = H, alkyl, Ac; R1 = H, halo, cyano, NO2, CF3, OCF3, alkyl, OH, alkylcarbonyloxy, alkoxy; R2-R5 = H, (halo)alkyl; R6 = H, halo, (halo)alkyl, cyano, alkoxy, aminocarbonyl, carboxy, alkylcarbonyl, (halo)alkoxy; R7 = H, halo, (halo)alkyl, dialkylaminoalkylaminocarbonyl, alkoxy, aminocarbonyl, ureido, aminosulfonyl, alkylsulfonylaminoalkylamino, aminosulfonylamino, heteroaryloxy, ureidoalkylaminocarbonyl, etc.; ≥1 of R2-R5 = alkyl], were prepared. Thus, 2-(2-amino-4-chlorophenoxy)-1-[4-(4-fluorophenoxy)piperidin-1-yl]ethanone (preparation given) in CH2Cl2 was treated with Et3N and Ph chloroformate. The reaction was stirred at ambient temperature for 4 h, concentrated in vacuo, and the resulting residue dissolved in methanol followed by bubbling in ammonia gas for 10 min and stirred overnight at ambient temperature to give [5-chloro-2-[2-[4-(4-fluorophenoxy)piperidin-1-yl]-2-oxoethoxy]phenyl]urea. I inhibited chemotaxis with IC50 <10 μM.

TI Preparation of piperidinylketones as selective inhibitors of macrophage inflammatory protein 1α (MIP-1α) binding to CCR1 chemokine receptors.

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

PRAI US 2002-397108P P 20020718 <--  
WO 2003-IB2876 W 20030707

L3 ANSWER 8 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2004:41226 CAPLUS [Full-text](#)

DOCUMENT NUMBER: 140:105321

TITLE: Methods and compositions relating to isoleucine

boroproline compounds

INVENTOR(S): Adams, Sharlene; Miller, Glenn T.; Jesson, Michael I.;

Jones, Barry

PATENT ASSIGNEE(S): Point Therapeutics, Inc., USA

SOURCE: PCT Int. Appl., 152 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

## PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004004658	A2	20040115	WO 2003-US21405	
20030709 <---				
WO 2004004658	A3	20050804		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZW				
AZ, BY, EE, ES, SK, TR, TD, TG				
CA 2491466	A1	20040115	CA 2003-2491466	
20030709 <---				
AU 2003265264	A1	20040123	AU 2003-265264	
20030709 <---				
US 20040077601	A1	20040422	US 2003-616694	
20030709 <---				
US 20050084490	A1	20050421	US 2003-616409	
20030709 <---				
EP 1578434	A2	20050928	EP 2003-763380	
20030709 <---				
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
JP 2006507352	T	20060302	JP 2004-562634	
20030709 <---				
CN 1802090	A	20060712	CN 2003-821282	
20030709 <---				
CN 1826129	A	20060830	CN 2003-821281	
20030709 <---				
IN 2005KN00151	A	20050916	IN 2005-KN151	
20050208 <---				
PRIORITY APPLN. INFO.:			US 2002-394856P	P
20020709 <---				
20021001 <---			US 2002-414978P	P
			US 2003-466435P	P
20030428			WO 2003-US21405	W
20030709				
OTHER SOURCE(S):	MARPAT 140:105321			

AB A method for treating subjects with, inter alia, abnormal cell proliferation or infectious disease using agents of formula (I,  $\text{AmNHCH}(\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3)\text{COA1R}$ ) (where Am and A1 are amino acids and R = organo boronates, organo phosphonates, fluoroalkyl ketones, aliphatic ketones, N-peptidyl-O-(acylhydroxylamines), azapeptides, azetidines, fluoroolefins dipeptide isosteres, peptidyl ( $\alpha$ -aminoalkyl) phosphonate esters, aminoacyl pyrrolidine-2-nitriles and 4-cyanothiazolidines) is claimed. Methods for stimulating an immune response using the compounds of the invention are also claimed. Compounds containing Ile-boroPro compounds are also provided as are kits containing the compounds. The invention embraces the use of these compounds alone or in combination with other therapeutic agents.

TI Methods and compositions relating to isoleucine boroproline compounds

REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

PRAI US 2002-394856P	P	20020709	<--
US 2002-414978P	P	20021001	<--
US 2003-466435P	P	20030428	
WO 2003-US21405	W	20030709	

L3 ANSWER 9 OF 23 CAPLUS COPYRIGHT 2009 ACS ON STN

ACCESSION NUMBER: 2003:836762 CAPLUS Full-text

DOCUMENT NUMBER: 139:350474

TITLE: Preparation and compositions of nitrosothio (hetero)cyclic nitric oxide donors

INVENTOR(S): Fang, Xinqin; Garvey, David S.; Gaston, Ricky D.; Lin,

Chia-en; Ranatunga, Ramani R.; Richardson,

Stewart K.;

Wang, Tiansheng; Wang, Weiheng; Wey, Shioh-jyi Nitromed, Inc., USA

PATENT ASSIGNEE(S):

SOURCE: PCT Int. Appl., 138 pp.

CODEN: PIXXD2

DOCUMENT TYPE:

Patent

LANGUAGE:



English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	----
----				
WO 2003086282	A2	20031023	WO 2003-US10562	
20030407 <--				
WO 2003086282	A3	20040429		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,				
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD,				
GE, GH,				
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,				
LK, LR,				
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ,				

OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR,  
 TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW  
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM,  
 AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,  
 EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI,  
 SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN,  
 TD, TG CA 2480832 A1 20031023 CA 2003-2480832  
 20030407 <-- AU 2003223491 A1 20031027 AU 2003-223491  
 20030407 <-- US 20030203915 A1 20031030 US 2003-407420  
 20030407 <-- EP 1497268 A2 20050119 EP 2003-719621  
 20030407 <-- R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,  
 MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK  
 JP 2005537223 T 20051208 JP 2003-583309  
 20030407 <-- PRIORITY APPLN. INFO.: US 2002-369873P P  
 20020405 <-- WO 2003-US10562 W  
 20030407  
 OTHER SOURCE(S): MARPAT 139:350474  
 GI



AB Title compds. I [wherein U = O, S, or NRaRi; V = NO or NO2; X9 = CR10 or N; Y9 = CR6R7, NRi, NR25, NRiCR6R7, CR6R7NRi, CR2R3CR6R7, or CR6R7CR2R3; Y10 = CR8R9 or CR8R9CR17R18; R2-R9, R17, and R18 = independently H or alkyl; or R2R3, R4R5, R6R7, or R8R9 = independently oxo; or R4 and R7 together with the C's to which they are attached = cycloalkyl; or CR6R7 = cycloalkyl; R6 and R9 taken together with the C's to which they are attached =



(bridged)cycloalkyl, heterocyclyl, or aryl with the proviso that R7 and R8 are not present; R4 and R25 taken together with the C and N to which they are attached = heterocyclyl; Ra = lone pair of electrons, H, or (aryl)alkyl; Re and Rf = independently H, halo, OH, or (un)substituted (cyclo)alkyl, heterocyclyl, alkoxy, amino, aryl, etc.; or CReRf = heterocyclyl or (bridged) cycloalkyl; Ri = H or (un)substituted alkyl, aryl, carboxamido, sulfonamido, etc.; n = 0-3; and pharmaceutically acceptable salts thereof] were prepared as novel nitric oxide donors for use in compns. comprising at least one nitric oxide donor and optionally at least one therapeutic agent. The nitric oxide donors donate, transfer or release nitric oxide, and/or elevate endogenous levels of endothelium-derived relaxing factor, and/or stimulate endogenous synthesis of nitric oxide and/or are substrates for nitric oxide synthase and are capable of releasing nitric oxide or indirectly delivering or transferring nitric oxide to targeted sites under physiol. conditions (no data). For example, 2-[2-(nitrosothio)adamantan-2-yl]acetic acid was esterified with 3-nitrooxy-2,2-bis(nitrooxymethyl)propan-1-ol in the presence of 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide•HCl and 4-dimethylaminopyridine in CH2Cl2 to give II (18%). The latter inhibited proliferation of human coronary artery smooth muscle cells with IC50 of 5 µM. In general, the nitrosylated compds. tested in this assay inhibited proliferation of vascular smooth muscle cells, while the corresponding non-nitrosylated derivs. showed no inhibition, slight inhibition, or exhibited much higher IC50 values. Thus, the invention provides methods for treating cardiovascular diseases, for the inhibition of platelet aggregation and platelet adhesion caused by the exposure of blood to a medical device, for treating pathol. conditions resulting from abnormal cell proliferation, transplantation rejections, autoimmune, inflammatory, proliferative, hyperproliferative, vascular diseases, for reducing scar tissue or for inhibiting wound contraction, particularly the prophylactic and/or therapeutic treatment of restenosis (no data). The invention also provides methods for treating inflammation, pain, fever, gastrointestinal disorders, respiratory disorders, and sexual dysfunctions (no data). In addition, the invention provides novel compns. and kits comprising at least one nitric oxide donor and/or at least one therapeutic agent.

TI Preparation and compositions of nitrosothio (hetero)cyclic nitric oxide donors

REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

PRAI US 2002-369873P P 20020405 <--  
WO 2003-US10562 W 20030407

AB Title compds. I [wherein U = O, S, or NRaRi; V = NO or NO2; X9 = CR10 or N; Y9 = CR6R7, NRi, NR25, NRiCR6R7, CR6R7NRi, CR2R3CR6R7, or CR6R7CR2R3; Y10 = CR8R9 or CR8R9CR17R18; R2-R9, R17, and R18 = independently H or alkyl; or R2R3, R4R5, R6R7, or R8R9 = independently oxo; or R4 and R7 together with the C's to which they are attached = cycloalkyl; or CR6R7 = cycloalkyl; R6 and R9 taken together with the C's to which they are attached =

(bridged)cycloalkyl, heterocyclyl, or aryl with the proviso that R7 and R8 are not present; R4 and R25 taken together with the C and N to which they are attached = heterocyclyl; Ra = lone pair of electrons, H, or (aryl)alkyl; Re and Rf = independently H, halo, OH, or (un)substituted (cyclo)alkyl, heterocyclyl, alkoxy, amino, aryl, etc.; or CReRf = heterocyclyl or (bridged) cycloalkyl; Ri = H or (un)substituted alkyl, aryl, carboxamido, sulfonamido, etc.; n = 0-3; and pharmaceutically acceptable salts thereof] were prepared as novel nitric oxide donors for use in compns. comprising at least one nitric oxide donor and optionally at least one therapeutic agent. The nitric oxide donors donate, transfer or release nitric oxide, and/or elevate endogenous levels of endothelium-derived relaxing factor, and/or stimulate endogenous synthesis of nitric oxide and/or are substrates for nitric oxide synthase and are capable of releasing nitric oxide or indirectly delivering or transferring nitric oxide to targeted sites under physiol. conditions (no data). For example, 2-[2-(nitrosothio)adamantan-2-yl]acetic acid was esterified with 3-nitrooxy-2,2-bis(nitrooxymethyl)propan-1-ol in the presence of 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide•HCl and 4-dimethylaminopyridine in CH2Cl2 to give II (18%). The latter inhibited proliferation of human coronary artery smooth muscle cells with IC50 of 5 µM. In general, the nitrosylated compds. tested in this assay inhibited proliferation of vascular smooth muscle cells, while the corresponding non-nitrosylated derivs. showed no inhibition, slight inhibition, or exhibited much higher IC50 values. Thus, the invention provides methods for treating cardiovascular diseases, for the inhibition of platelet aggregation and platelet adhesion caused by the exposure of blood to a medical device, for treating pathol. conditions resulting from abnormal cell proliferation, transplantation rejections, autoimmune, inflammatory, proliferative, hyperproliferative, vascular diseases, for reducing scar tissue or for inhibiting wound contraction, particularly the prophylactic and/or therapeutic treatment of restenosis (no data). The invention also provides methods for treating inflammation, pain, fever, gastrointestinal disorders, respiratory disorders, and sexual dysfunctions (no data). In addition, the invention provides novel compns. and kits comprising at least one nitric oxide donor and/or at least one therapeutic agent.

L3 ANSWER 10 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 2003:777485 CAPLUS Full-text  
 DOCUMENT NUMBER: 139:272356  
 TITLE: Fungicidal compositions containing  
 benzamides in combination with other  
 fungicides  
 INVENTOR(S): Walker, Michael Paul; Foor, Stephen Ray  
 PATENT ASSIGNEE(S): E. I. Du Pont de Nemours & Co., USA; Walker,  
 Susannah  
 H. F.  
 SOURCE: PCT Int. Appl., 59 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
----- WO 2003079788	A2	20031002	WO 2003-US8205	
20030318 <--				
WO 2003079788	A3	20040219		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,				
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,				
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,				
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM,				
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT,				
TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,				
KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES,				
FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR,				
BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2003220361	A1	20031008	AU 2003-220361	
20030318 <--				
EP 1484970	A2	20041215	EP 2003-716661	
20030318 <--				
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
BR 2003008457	A	20050111	BR 2003-8457	
20030318 <--				
JP 2005520839	T	20050714	JP 2003-577631	
20030318 <--				
CN 1642421	A	20050720	CN 2003-806454	
20030318 <--				
ZA 2004005644	A	20050811	ZA 2004-5644	
20030318 <--				
RU 2314690	C2	20080120	RU 2004-130840	
20030318 <--				
IN 2004DN01996	A	20070406	IN 2004-DN1996	
20040612 <--				
US 20050164999	A1	20050728	US 2004-501126	
20040709 <--				
MX 2004009001	A	20041207	MX 2004-9001	
20040915 <--				
PRIORITY APPLN. INFO.:			US 2002-365764P	P
20020319 <--				
			WO 2003-US8205	W
20030318				
OTHER SOURCE(S):	MARPAT 139:272356			

AB Comps. for controlling plant diseases caused by fungal plant pathogens comprise: (a) a fungicidally effective amount of a compound (A) (R1) (R2) -N(R3)-W-B, or N-oxides, and agriculturally suitable salts thereof (A = substituted pyridinyl; B = substituted phenyl; W = C:O, C:S, or SO<sub>n</sub>; n = 1, 2; R1, R2 = H, or (un)substituted C1-C6 alkyl, C2-C6 alkenyl, C2-C6 alkynyl, or C3-C6 cycloalkyl; R3 = H, C1-C6 alkyl, C2-C6 alkenyl, C2-C6 alkynyl, or C3-C6 cycloalkyl, C2-C10 alkoxyalkyl, C2-C6 alkylcarbonyl, C2-C6 alkoxy carbonyl, C2-C6 alkylaminocarbonyl, or C3-C8 dialkylaminocarbonyl), and (b) at least one compound selected from the group consisting of (b1) alkylenebis(dithiocarbamate) fungicides; (b2) compds. acting at the bcl complex of the fungal mitochondrial respiratory electron transfer site; (b3) cymoxanil; (b4) compds. acting at the demethylase enzyme of the sterol biosynthesis pathway; (b5) morpholine and piperidine compds. that act on the sterol biosynthesis pathway; (b6) phenylamide fungicides; (b7) pyrimidinone fungicides; (b8) phthalimides; and (b9) fosetyl-aluminum.

TI Fungicidal compositions containing benzamides in combination with other fungicides

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

TI Fungicidal compositions containing benzamides in combination with other fungicides

PRAI US 2002-365764P P 20020319 <--  
WO 2003-US8205 W 20030318

=> d l3 ibib abs 11-20 ti hit

L3 ANSWER 11 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2003:757679 CAPLUS Full-text

DOCUMENT NUMBER: 139:276825

TITLE: Preparation of 8-arylquinoline PDE4 inhibitors

INVENTOR(S): Gallant, Michel; Lacombe, Patrick; Dube, Daniel;

Deschenes, Denis; MacDonald, Dwight; Dube,

Laurence

PATENT ASSIGNEE(S): Merck Frosst Canada & Co., Can.

SOURCE: PCT Int. Appl., 184 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent



LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003078397	A1	20030925	WO 2003-CA374	
20030317 <--				
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,			
	CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD,			

GE, GH,  
 LR, LS,  
 OM, PH,  
 TT, TZ,  
 RW: UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW  
 AZ, BY, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM,  
 EE, ES, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,  
 SK, TR, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI,  
 TD, TG BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN,  
 CA 2479069 A1 20030925 CA 2003-2479069  
 20030317 <--  
 AU 2003209896 A1 20030929 AU 2003-209896  
 20030317 <--  
 EP 1487797 A1 20041222 EP 2003-744288  
 20030317 <--  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,  
 MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK  
 US 20050245513 A1 20051103 US 2004-508261  
 20040917 <--  
 US 7144896 B2 20061205  
 PRIORITY APPLN. INFO.: US 2002-365088P P  
 20020318 <--  
 WO 2003-CA374 W  
 20030317  
 OTHER SOURCE(S): MARPAT 139:276825  
 GI



AB Title compds. I {wherein R1 = H, halo, or (un)substituted alkanoyl, (cyclo)alkyl, alkenyl, alkoxy, (hetero)aryl, CN, heterocycloalkyl, carbamoyl, sulfamoyl, etc.; R2 = H, halo, OH, or (un)substituted alkyl or alkoxy; R3 = absent or H, CO2H, or


(un)substituted (cycloalkyl)alkyl, alkanoyl, benzoyl, carbamoyl, etc.; R4 = (un)substituted Ph, pyrazolopyrimidinyl, benzothiazolyl, quinazolinyl, or heteroaryl; R5 = absent or H; R6 = absent, H, or alkyl; R7 = absent or H; X = O, S, N, C, or CO; wherein when X = O, S, or CO, then R6 and R7 are absent and when X = N, then R7 is absent; Y = C, S, N, SO2, O, or CO; wherein when Y = S, SO2, O, or CO, then R3 and R5 are absent and when Y = N, then R5 is absent; and pharmaceutically acceptable salts thereof] were prepared as phosphodiesterase IV (PDE4) inhibitors. For example, 3-(6-isopropylquinolin-8-yl)phenol was coupled with 1-chloromethyl-4-methanesulfonylbenzene in acetone to give II. One hundred sixteen invention compds. suppressed PDE4 with IC50 values ranging from 80  $\mu$ M to 0.029  $\mu$ M in assays evaluating LPS- and FMLP-induced inhibition of tumor necrosis factor  $\alpha$  (TNF- $\alpha$ ) and leukotriene B4 (LTB4) in human whole blood. In a test measuring IgE-mediated allergic pulmonary inflammation induced by inhalation of antigen by sensitized guinea pigs, administration of I resulted in a significant reduction in the eosinophilia and the accumulation of other inflammatory leukocytes and effected less inflammatory lung damage. One hundred forty-one invention compds. also inhibited the hydrolysis of cAMP to AMP by human recombinant phosphodiesterase IVa with IC50 values ranging from 150 nM to 0.056 nM. Thus, I and their pharmaceutical compns. are useful for the treatment or prevention of a variety of allergic, inflammatory, CNS, and other conditions (no data).

TI Preparation of 8-arylquinoline PDE4 inhibitors  
REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE  
FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE  
RE FORMAT  
PRAI US 2002-365088P P 20020318 <--  
WO 2003-CA374 W 20030317

L3 ANSWER 12 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 2003:656587 CAPLUS Full-text  
DOCUMENT NUMBER: 139:197374  
TITLE: Preparation of nicotinamides useful as PDE4  
inhibitors  
for treating diseases including inflammatory,  
allergic  
and respiratory diseases  
INVENTOR(S): Bailey, Simon; Gautier, Elisabeth Colette  
Louise;  
Henderson, Alan John; Magee, Thomas Victor;  
Marfat,  
Anthony; Mathias, John Paul; McLeod, Dale  
Gordon;  
Monaghan, Sandra Marina; Stammen, Blanda Luzia  
Christa  
PATENT ASSIGNEE(S): Pfizer Limited, UK; Pfizer Inc.  
SOURCE: PCT Int. Appl., 266 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 2  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----				
WO 2003068235	A1	20030821	WO 2003-IB439	
20030203 <--				
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CA 2475712	A1	20030821	CA 2003-2475712	
20030203 <--				
AU 2003245711	A1	20030904	AU 2003-245711	
20030203 <--				
EP 1476158	A1	20041117	EP 2003-739392	
20030203 <--				
EP 1476158	B1	20071114		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
BR 2003007564	A	20041221	BR 2003-7564	
20030203 <--				
JP 2005522450	T	20050728	JP 2003-567417	
20030203 <--				
CN 1652782	A	20050810	CN 2003-808186	
20030203 <--				
NZ 534197	A	20070126	NZ 2003-534197	
20030203 <--				
AT 378049	T	20071115	AT 2003-739392	
20030203 <--				
ES 2292988	T3	20080316	ES 2003-739392	
20030203 <--				
US 20030220361	A1	20031127	US 2003-360100	
20030206 <--				
US 20030220366	A1	20031127	US 2003-361062	
20030206 <--				
US 6949573	B2	20050927		
US 20040224975	A1	20041111	US 2004-865263	
20040609 <--				
US 7060717	B2	20060613		
IN 2004DN02070	A	20050401	IN 2004-DN2070	
20040719 <--				
MX 2004007737	A	20041015	MX 2004-7737	

20040810 <--				
NO 2004003793	A	20041021	NO 2004-3793	
20040910 <--				
US 20060014780	A1	20060119	US 2005-229395	
20050916 <--				
ZA 2004005803	A	20060531	ZA 2004-5803	
20060316 <--				
PRIORITY APPLN. INFO.:			GB 2002-3196	A
20020211 <--				
			GB 2002-20999	A
20020910 <--				
			GB 2002-24453	A
20021021 <--				
			GB 2002-27139	A
20021120 <--				
			US 2002-361991P	P
20020305 <--				
			GB 2002-20984	A
20020910 <--				
			US 2002-414247P	P
20020926 <--				
			US 2002-414304P	P
20020926 <--				
			GB 2002-24454	A
20021021 <--				
			US 2002-425406P	P
20021112 <--				
			US 2002-425474P	P
20021112 <--				
			GB 2002-27140	A
20021120 <--				
			US 2002-433330P	P
20021213 <--				
			US 2002-433336P	P
20021213 <--				
			WO 2003-IB439	W
20030203				
			US 2003-361062	A3
20030206				
			US 2004-865263	A1
20040609				
OTHER SOURCE(S) :	MARPAT 139:197374			
GI				



I

AB The invention relates to nicotinamides (shown as I; variables defined below; e.g. anti-2-(benzo[1,3]dioxol-5-yloxy)-N-[4-(2-hydroxybenzoylamino)cyclohexyl]nicotinamide) and to processes for the preparation of, intermediates used in the preparation of,




comps. containing and the uses of, such derivs. The nicotinamide derivs. according to the present invention are phosphodiesterase-4 inhibitors and are useful in numerous diseases, disorders and conditions, in particular inflammatory, allergic, respiratory diseases, disorders and conditions, as well as wounds. For I: R1 and R2 = H, halo, cyano, (C1-C4)alkyl and (C1-C4)alkoxy; X is -O-, -S- or -NH-; R3 = Ph, naphthyl, heteroaryl and (C3-C8)cycloalkyl or the bicyclic groups benzodioxol-5-yl, benzofuran-5-yl, benzofuran-6-yl, indan-5-yl; Y = 4-HNcyclohexyl, piperidin-1,4-diyl, 8-azabicyclo[3.2.1]octane-3,8-diyl, and 4-R5Ncyclohexyl wherein in each the N is bonded to Z in I and R5 = (C1-C4)alkyl and phenyl(C1-C4)alkyl. Z = C(O), C(O)NH, SO2, SO2NH, C(O)CH2NHSO2, SO2NHC(O), C(O)CH2NHC(O) wherein the left end is bonded to Y and the other end to R4; or alternatively Y-Z together = 4-NHC(O)cyclohexyl; R4 = Ph, naphthyl heteroaryl and (C3-C8)cycloalkyl, (un)substituted (C1-C6)alkyl; addnl. details including provisos are given in the claims. The antiinflammatory properties of 72 examples of I are demonstrated by their ability to inhibit TNF $\alpha$  release from human peripheral blood mononuclear cells, e.g. IC50 = 0.014 nM for syn-2-(3,4-difluorophenoxy)-5-fluoro-N-[4-(2-hydroxy-5-methylbenzoylamino)cyclohexyl]nicotinamide. About 200 example preps. of I and 75 of intermediates are included. For example, to prepare anti-2-[(benzo[1,3]dioxol-5-yl)oxy]-N-[4-[(2-hydroxybenzoyl)amino]cyclohexyl]nicotinamide (160.7 mg), 2-hydroxybenzoic acid (0.767 mmol), 1-hydroxybenzotriazole hydrate (1.15 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.15 mmol) were stirred in DMF (5 mL) under an atmospheric of N2 at room temperature for 1.5 h. Anti-N-(4-aminocyclohexyl)-2-[(benzo[1,3]dioxol-5-yl)oxy]nicotinamide hydrochloride (0.767 mmol; preparation given) and N-methylmorpholine (0.767 mmol) were then added, and the reaction mixture stirred at room temperature for a further 18 h.

L3 ANSWER 13 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 2002:695687 CAPLUS Full-text  
 DOCUMENT NUMBER: 137:212298  
 TITLE: Fungicidal compositions based on  
 pyridylmethylbenzamide derivatives and complex  
 III inhibiting compounds  
 INVENTOR(S): Holah, David Stanley; Dancer, Jane Elisabeth;  
 Latorse, Marie-Pascale; Mercer, Richard  
 PATENT ASSIGNEE(S): Aventis CropScience SA, Fr.  
 SOURCE: PCT Int. Appl., 44 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002069712	A1	20020912	WO 2002-EP4613	

20020307 <--

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,  
 CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD,  
 GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
 LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ,  
 OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR,  
 TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW  
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT,  
 BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT,  
 SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN,  
 TD, TG  
 FR 2821718 A1 20020913 FR 2001-3140  
 20010308 <--  
 FR 2821718 B1 20030613  
 EG 23127 A 20040428 EG 2002-239  
 20020305 <--  
 AU 2002304701 A1 20020919 AU 2002-304701  
 20020307 <--  
 EP 1365653 A1 20031203 EP 2002-732676  
 20020307 <--  
 EP 1365653 B1 20041201  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,  
 MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR  
 JP 2004523564 T 20040805 JP 2002-568906  
 20020307 <--  
 AT 283630 T 20041215 AT 2002-732676  
 20020307 <--  
 PT 1365653 T 20050331 PT 2002-732676  
 20020307 <--  
 ES 2231700 T3 20050516 ES 2002-732676  
 20020307 <--  
 US 20040106578 A1 20040603 US 2003-471124  
 20030908 <--  
 KR 838538 B1 20080617 KR 2003-711828  
 20030908 <--  
 US 20070293549 A1 20071220 US 2007-879592  
 20070718 <--  
 PRIORITY APPLN. INFO.: FR 2001-3140 A  
 20010308 <-- WO 2002-EP4613 W  
 20020307 <-- US 2003-471124 B3  
 20030908  
 OTHER SOURCE(S): MARPAT 137:212298  
 GI



AB A fungicide compns. comprises (a) a pyridylmethylbenzamide derivative I (Markush included), and (b) at least one compound capable of inhibiting the transport of electrons of the respiratory chain of mitochondrial ubiquinol-ferricytochrome-c oxidoreductase or complex III in phytopathogenic fungal organisms. The composition is used as preventive or curative agent for fighting against phytopathogenic fungi of crops by applying on the aerial parts of plants an efficient and non-phytotoxic amount of said fungicide compns.

TI Fungicidal compositions based on pyridylmethylbenzamide derivatives and complex III inhibiting compounds

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

TI Fungicidal compositions based on pyridylmethylbenzamide derivatives and complex III inhibiting compounds

PI	WO 2002069712 A1	20020912			
	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
----	-----	----	-----	-----	----
PI	WO 2002069712	A1	20020912	WO 2002-EP4613	
20020307 <--					
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW				
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
FR 2821718	A1	20020913	FR 2001-3140		
20010308 <--					
FR 2821718	B1	20030613			
EG 23127	A	20040428	EG 2002-239		
20020305 <--					
AU 2002304701	A1	20020919	AU 2002-304701		
20020307 <--					

EP 1365653 A1 20031203 EP 2002-732676  
 20020307 <--  
 EP 1365653 B1 20041201  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,  
 MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR  
 JP 2004523564 T 20040805 JP 2002-568906  
 20020307 <--  
 AT 283630 T 20041215 AT 2002-732676  
 20020307 <--  
 PT 1365653 T 20050331 PT 2002-732676  
 20020307 <--  
 ES 2231700 T3 20050516 ES 2002-732676  
 20020307 <--  
 US 20040106578 A1 20040603 US 2003-471124  
 20030908 <--  
 KR 838538 B1 20080617 KR 2003-711828  
 20030908 <--  
 US 20070293549 A1 20071220 US 2007-879592  
 20070718 <--  
 PR AI FR 2001-3140 A 20010308 <--  
 WO 2002-EP4613 W 20020307 <--  
 US 2003-471124 B3 20030908

L3 ANSWER 14 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 2002:521462 CAPLUS Full-text  
 DOCUMENT NUMBER: 137:88442  
 TITLE: Incensole and furanogermacrene and compounds  
 in

treatment for inhibiting neoplastic lesions

and  
 microorganisms  
 INVENTOR(S): Shanahan-Pendergast, Elisabeth  
 PATENT ASSIGNEE(S): Ire.  
 SOURCE: PCT Int. Appl., 68 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	----
WO 2002053138	A2	20020711	WO 2002-IE1	
20020102 <--				
WO 2002053138	A3	20020919		
W: AE, AG, AT, AU, BB, BG, CA, CH, CN, CO, CU, CZ, LU, LV,				
MA, MD, UA, UG, US, VN, YU, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, AT, BE, CH, CY, DE,				
ES, FI, ML, MR, NE, SN, TD, TG				
AU 2002219472	A1	20020716	AU 2002-219472	
20020102 <--				
EP 1351678	A2	20031015	EP 2002-727007	

20020102 <--  
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,  
MC, PT,  
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR  
US 20040092583 A1 20040513 US 2004-250535  
20040102 <--  
PRIORITY APPLN. INFO.: IE 2001-2 A  
20010102 <--  
WO 2002-IE1 W  
20020102 <--  
OTHER SOURCE(S): MARPAT 137:88442  
AB The invention discloses the use of incensole and/or  
furanogermacrene, derivs. metabolites and precursors thereof in  
the treatment of neoplasia, particularly resistant neoplasia and  
immunodysregulatory disorders. These compds. can be administered  
alone or in combination with conventional chemotherapeutic,  
antiviral, antiparasite agents, radiation and/or surgery.  
Incensole and furanogermacrene and their mixture showed antitumor  
activity against various human carcinomas and melanomas and  
antimicrobial activity against Staphylococcus aureus and  
Enterococcus faecalis.  
TI Incensole and furanogermacrene and compounds in treatment for  
inhibiting  
neoplastic lesions and microorganisms  
REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE  
FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE  
RE FORMAT  
PI WO 2002053138 A2 20020711  
PATENT NO. KIND DATE APPLICATION NO. DATE  
-----  
----  
PI WO 2002053138 A2 20020711 WO 2002-IE1  
20020102 <--  
WO 2002053138 A3 20020919  
W: AE, AG, AT, AU, BB, BG, CA, CH, CN, CO, CU, CZ, LU, LV,  
MA, MD,  
UA, UG, US, VN, YU, RU, TJ, TM  
RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, AT, BE, CH, CY, DE,  
ES, FI,  
ML, MR, NE, SN, TD, TG  
AU 2002219472 A1 20020716 AU 2002-219472  
20020102 <--  
EP 1351678 A2 20031015 EP 2002-727007  
20020102 <--  
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,  
MC, PT,  
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR  
US 20040092583 A1 20040513 US 2004-250535  
20040102 <--  
PRAI IE 2001-2 A 20010102 <--  
WO 2002-IE1 W 20020102 <--  
L3 ANSWER 15 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 2002:314940 CAPLUS Full-text  
DOCUMENT NUMBER: 136:340711

TITLE: Bridged piperazine derivatives, specifically  
3,8-diazabicyclo[3.2.1]octane,  
8-azabicyclo[3.2.1]octane,  
2,5-diazabicyclo[2.2.2]octane, and  
3,9-diazabicyclo[3.3.1]nonane derivatives,  
useful as  
receptors, inhibitors of chemokines binding to CCR1  
for treating inflammation and other immune  
disorders.  
INVENTOR(S): Blumberg, Laura Cook; Brown, Matthew Frank;  
Glaude,  
Ronald Paul; Poss, Christopher Stanley  
PATENT ASSIGNEE(S): Pfizer Products Inc., USA  
SOURCE: PCT Int. Appl., 89 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
----	-----	----	-----	-----	-----
20011004 <--	WO 2002032901	A2	20020425	WO 2001-IB1844	
	WO 2002032901	A3	20020725		
CH, CN,	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,				
GE, GH,	CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD,				
LK, LR,	GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,				
PH, PL,	LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ,				
UA, UG,	PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ,				
	US, UZ, VN, YU, ZA, ZW				
CH, CY,	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE,				
TR, BF,	DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE,				
	BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
20011004 <--	CA 2423789	A1	20020425	CA 2001-2423789	
	AU 2001092160	A	20020429	AU 2001-92160	
20011004 <--	EP 1326867	A2	20030716	EP 2001-972389	
20011004 <--	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,				
MC, PT,	IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
	EE 200300189	A	20031015	EE 2003-189	
20011004 <--	BR 2001014697	A	20031118	BR 2001-14697	
20011004 <--	HU 2003001442	A2	20031229	HU 2003-1442	

20011004 <--  
     HU 2003001442                   A3     20070328  
     JP 2004511558                   T     20040415     JP 2002-536283  
 20011004 <--  
     NZ 524742                       A     20041224     NZ 2001-524742  
 20011004 <--  
     US 20020119961                  A1     20020829     US 2001-972177  
 20011005 <--  
     IN 2003MN00309                  A     20050211     IN 2003-MN309  
 20030317 <--  
     ZA 2003002157                  A     20040422     ZA 2003-2157  
 20030318 <--  
     BG 107655                       A     20040130     BG 2003-107655  
 20030320 <--  
     NO 2003001572                  A     20030610     NO 2003-1572  
 20030408 <--  
     MX 2003003475                  A     20030714     MX 2003-3475  
 20030416 <--  
 PRIORITY APPLN. INFO.:                               US 2000-241804P     P  
 20001019 <--   WO 2001-1B1844     W  
  
 20011004 <--  
 OTHER SOURCE(S):                   MARPAT 136:340711  
 GI



AB    Comps. I and their pharmaceutically acceptable salts, useful for treatment of inflammation and other immune disorders, are disclosed [wherein: n = 1-5; m = 1-5; q = 0-1; a, b, c = (CH<sub>2</sub>)<sub>0-4</sub> (independently); a, b, and c cannot all be null; if a and/or c is not null, then b must be null; W = CH or N; X = CO, C(S), or CH<sub>2</sub>; Y = CH<sub>2</sub>; Z = O, (un)substituted NH or (un)substituted CH<sub>2</sub>; R = certain (un)substituted (hetero)aryl or (hetero)cycloalkyl; R<sub>1</sub> = (independently) H, OH, SO<sub>3</sub>H, halo, alkyl, SH, CF<sub>3</sub>, wide variety of other substituents]. The comps. are useful for treatment of a wide variety of diseases and disorders, which are cited specifically in claims. Approx. 100 specific examples of I are given, many with synthetic details. For example, 3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]octan-2-one (preparation

given) underwent a sequence of: (1) reduction of the amide carbonyl using LiAlH<sub>4</sub> (94%); (2) 8-N-acylation with chloroacetyl chloride (69%); and (3) etherification with 2-nitro-4-trifluoromethylphenol (58%), to give title compound II. In a bioassay for the ability to inhibit chemotaxis of various cells (THP-1 cells, primary human monocytes, or primary lymphocytes) in vitro, all example compds. had IC<sub>50</sub> values of less than 10 μM.

TI Bridged piperazine derivatives, specifically 3,8-diazabicyclo[3.2.1]octane, 8-azabicyclo[3.2.1]octane, 2,5-diazabicyclo[2.2.2]octane, and 3,9-diazabicyclo[3.3.1]nonane derivatives, useful as inhibitors of chemokines binding to CCRL1 receptors,

for treating inflammation and other immune disorders.

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

# RE FORMAT

PI	WO 2002032901 A2	20020425			
	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
----	-----	----	-----	-----	----
PI	WO 2002032901	A2	20020425	WO 2001-IB1844	
20011004 <--					
	WO 2002032901	A3	20020725		
CH, CN,	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,				
GE, GH,	CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD,				
LK, LR,	GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,				
PH, PL,	LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ,				
UA, UG,	PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ,				
	US, UZ, VN, YU, ZA, ZW				
CH, CY,	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE,				
TR, BF,	DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE,				
	BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	CA 2423789	A1	20020425	CA 2001-2423789	
20011004 <--					
	AU 2001092160	A	20020429	AU 2001-92160	
20011004 <--					
	EP 1326867	A2	20030716	EP 2001-972389	
20011004 <--					
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,				
MC, PT,	IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
	EE 200300189	A	20031015	EE 2003-189	
20011004 <--					
	BR 2001014697	A	20031118	BR 2001-14697	
20011004 <--					
	HU 2003001442	A2	20031229	HU 2003-1442	
20011004 <--					
	HU 2003001442	A3	20070328		
	JP 2004511558	T	20040415	JP 2002-536283	



20011004 <--  
 NZ 524742 A 20041224 NZ 2001-524742  
 20011004 <--  
 US 20020119961 A1 20020829 US 2001-972177  
 20011005 <--  
 IN 2003MN00309 A 20050211 IN 2003-MN309  
 20030317 <--  
 ZA 2003002157 A 20040422 ZA 2003-2157  
 20030318 <--  
 BG 107655 A 20040130 BG 2003-107655  
 20030320 <--  
 NO 2003001572 A 20030610 NO 2003-1572  
 20030408 <--  
 MX 2003003475 A 20030714 MX 2003-3475  
 20030416 <--  
 PRAI US 2000-241804P P 20001019 <--

L3 ANSWER 16 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 2001:569687 CAPLUS Full-text  
 DOCUMENT NUMBER: 135:148587  
 TITLE: Synergistic fungicidal compositions  
 containing N-acetonylbenzamides  
 INVENTOR(S): Young, David Hamilton; Wilson, Willie Joe;  
 Egan, Anne  
 PATENT ASSIGNEE(S): Ritchie; Michelott, Enrique Luis  
 Rohm and Haas Company, USA  
 SOURCE: U.S., 8 pp., Cont.-in-part of U.S. 6,075,047.  
 CODEN: USXXAM  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 4  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	-----	-----	-----	-----
US 6270810	B1	20010807	US 2000-561842	
20000428 <--				
US 6004947	A	19991221	US 1998-148604	
19980904 <--				
US 6075047	A	20000613	US 1999-433676	
19991104 <--				
PRIORITY APPLN. INFO.:			US 1998-72725P	P
19980127 <--				
			US 1998-148604	A3
19980904 <--				
			US 1999-433676	A2
19991104 <--				
AB The title compns. comprise a N-acetonylbenzamide derivative and a 2nd compound from the group consisting of an inhibitor of respiration at cytochrome complex III, ziram, fluazinam, zarilamide, chlorothalonil, propamocarb, folpet, fosetyl-aluminum or a fungitoxic metabolite thereof, a triphenyltin type fungicide and a copper fungicide.				
TI Synergistic fungicidal compositions containing N-acetonylbenzamides				

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE  
FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

TI Synergistic fungicidal compositions containing N-  
acetylbenzamides

PI US 6270810 B1 20010807

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
------------	------	------	-----------------	------

-----	----	-----	-----	----
-------	------	-------	-------	------

PI	US 6270810	B1	20010807	US 2000-561842
----	------------	----	----------	----------------

20000428 <--				
--------------	--	--	--	--

US 6004947	A	19991221	US 1998-148604	
------------	---	----------	----------------	--

19980904 <--				
--------------	--	--	--	--

US 6075047	A	20000613	US 1999-433676	
------------	---	----------	----------------	--

19991104 <--				
--------------	--	--	--	--

PRAI	US 1998-72725P	P	19980127 <--	
------	----------------	---	--------------	--

US 1998-148604	A3	19980904 <--		
----------------	----	--------------	--	--

US 1999-433676	A2	19991104 <--		
----------------	----	--------------	--	--

AB The title comps. comprise a N-acetylbenzamide derivative and a 2nd compound from the group consisting of an inhibitor of respiration at cytochrome complex III, ziram, fluazinam, zarilamide, chlorothalonil, propamocarb, folpet, fosetyl-aluminum or a fungitoxic metabolite thereof, a triphenyltin type fungicide and a copper fungicide.

ST synergism fungicide acetylbenzamide deriv compn

IT Albugo  
Oomycetes  
Peronospora  
Phytophthora  
Plasmopara  
Pseudoperonospora  
(control with synergistic fungicidal comps. containing N-acetylbenzamides)

IT Fungicides  
(synergistic, agrochem.; comps. containing N-acetylbenzamides)

IT	238739-68-9	238739-69-0	238739-70-3	238739-71-4	238739-72-5
----	-------------	-------------	-------------	-------------	-------------

350482-25-6	352272-55-0				
-------------	-------------	--	--	--	--

RL: AGR (Agricultural use); BIOL (Biological study); USES (Uses)  
(synergistic fungicidal composition)

IT 156052-68-5D, mixts. containing  
RL: AGR (Agricultural use); BIOL (Biological study); USES (Uses)  
(synergistic fungicidal comps.)

L3 ANSWER 17 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2001:559978 CAPLUS Full-text

DOCUMENT NUMBER: 135:103785

TITLE: Synergistic fungicidal compositions  
containing N-acetylbenzamides

INVENTOR(S): Young, David Hamilton; Wilson, Willie Joe;  
Egan, Anne

PATENT ASSIGNEE(S): Ritchie; Michelott, Enrique Luis  
Rohm and Haas Co., USA

SOURCE: U.S., 8 pp., Cont.-in-part of U.S. 6,075,047.  
CODEN: USXXAM

DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 4  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
20000428 <--	B1	20010731	US 2000-561841	
19980904 <--	A	19991221	US 1998-148604	
19991104 <--	A	20000613	US 1999-433676	
PRIORITY APPLN. INFO.: 19980127 <--			US 1998-72725P	P
19980904 <--			US 1998-148604	A3
19991104 <--			US 1999-433676	A2

AB The title compns. active against phytopathogenic fungi comprise an N-acetonylbenzamide derivative, preferably N-[3'-(1'-chloro-3'-methyl-2'-oxopentane)]-3,5-dichloro-4- methylbenzamide, and a second fungicidally-active compound selected from respiration inhibitors at cytochrome complex III, ziram, fluazinam, zarilamide, chlorothalonil, propamocarb, folpet, fosetyl-aluminum or a fungitoxic metabolite thereof, a triphenyltin type fungicide or a copper containing fungicide .

TI Synergistic fungicidal compositions containing N-acetonylbenzamides

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

TI Synergistic fungicidal compositions containing N-acetonylbenzamides

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
20000428 <--	B1	20010731	US 2000-561841	
19980904 <--	A	19991221	US 1998-148604	
19991104 <--	A	20000613	US 1999-433676	
PRAI US 1998-72725P	P	19980127 <--		
US 1998-148604	A3	19980904 <--		
US 1999-433676	A2	19991104 <--		

AB The title compns. active against phytopathogenic fungi comprise an N-acetonylbenzamide derivative, preferably N-[3'-(1'-chloro-3'-methyl-2'-oxopentane)]-3,5-dichloro-4- methylbenzamide, and a second fungicidally-active compound selected from respiration inhibitors at cytochrome complex III, ziram, fluazinam, zarilamide, chlorothalonil, propamocarb, folpet, fosetyl-aluminum or a fungitoxic metabolite thereof, a triphenyltin type fungicide or a copper containing fungicide .

ST synergism fungicide agrochem acetonylbenzamide deriv  
 IT Albugo  
 Oomycetes  
 Peronospora  
 Phytophthora  
 Plasmopara  
 Pseudoperonospora  
 (control by synergistic fungicidal compns. containing N-acetonylbenzamides)  
 IT Fungicides  
 (synergistic, agrochem.; compns. containing N-acetonylbenzamides)  
 IT 238739-68-9 238739-69-0 238739-70-3 238739-71-4 238739-72-5  
 350482-24-5 350482-25-6  
 RL: AGR (Agricultural use); BIOL (Biological study); USES (Uses)  
 (synergistic fungicidal composition)  
 IT 156052-68-5D, mixts. containing  
 RL: AGR (Agricultural use); BIOL (Biological study); USES (Uses)  
 (synergistic fungicidal compns.)

L3 ANSWER 18 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 2001:537393 CAPLUS Full-text  
 DOCUMENT NUMBER: 135:103783  
 TITLE: Synergistic fungicidal compositions  
 containing a N-acetonylbenzamide derivative  
 INVENTOR(S): Young, David Hamilton; Wilson, Willie Joe;  
 Egan, Anne  
 PATENT ASSIGNEE(S): Ritchie, Michelott, Enrique Luis  
 Rohm and Haas Company, USA  
 SOURCE: U.S., 8 pp., Cont.-in-part of U.S. 6,075,047.  
 CODEN: USXXAM  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 4  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	----
US 6264993	B1	20010724	US 2000-561037	
20000428 <--				
US 6004947	A	19991221	US 1998-148604	
19980904 <--				
US 6075047	A	20000613	US 1999-433676	
19991104 <--				
PRIORITY APPLN. INFO.:			US 1998-72725P	P
19980127 <--				
			US 1998-148604	A3
19980904 <--				
			US 1999-433676	A2
19991104 <--				
OTHER SOURCE(S):	MARPAT 135:103783			

AB The invention relates to synergistic fungicidal compns. comprising N-[3'-(1'-chloro-3'-methyl-2'-oxopentane)]-3,5-dichloro-4-methylbenzamide and a second fungicidally-active compound selected from an inhibitor of respiration at cytochrome complex III, ziram,

fluazinam, zarilamide, chlorothalonil, propamocarb, folpet, fosetyl-aluminum or a fungitoxic metabolite thereof, phosphorous acid or a salt thereof, a triphenyltin type fungicide and a copper containing fungicide to plant seed, to plant foliage or to a plant growth medium.

TI Synergistic fungicidal compositions containing a N-acetylbenzamide derivative

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

TI Synergistic fungicidal compositions containing a N-acetylbenzamide derivative

PI US 6264993 B1 20010724

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	---	-----	-----	----

PI	US 6264993	B1	20010724	US 2000-561037
----	------------	----	----------	----------------

20000428 <--

US 6004947	A	19991221	US 1998-148604
------------	---	----------	----------------

19980904 <--

US 6075047	A	20000613	US 1999-433676
------------	---	----------	----------------

19991104 <--

PRAI	US 1998-72725P	P	19980127	<--
------	----------------	---	----------	-----

US 1998-148604	A3	19980904	<--
----------------	----	----------	-----

US 1999-433676	A2	19991104	<--
----------------	----	----------	-----

AB The invention relates to synergistic fungicidal compns. comprising N-[3'-(1'-chloro-3'-methyl-2'-oxopentane)]-3,5-dichloro-4-methylbenzamide and a second fungicidally-active compound selected from an inhibitor of respiration at cytochrome complex III, ziram, fluazinam, zarilamide, chlorothalonil, propamocarb, folpet, fosetyl-aluminum or a fungitoxic metabolite thereof, phosphorous acid or a salt thereof, a triphenyltin type fungicide and a copper containing fungicide to plant seed, to plant foliage or to a plant growth medium.

ST synergism fungicide compn acetylbenzamide deriv

IT Albugo

Peronospora

Phytophthora

Plasmopara

Pseudoperonospora

(synergistic fungicidal composition for control of)

IT Fungicides

(synergistic; containing a N-acetylbenzamide derivative)

IT	238739-68-9	238739-69-0	238739-70-3	238739-71-4	238739-72-5
----	-------------	-------------	-------------	-------------	-------------

350482-25-6 350482-49-4

RL: AGR (Agricultural use); BIOL (Biological study); USES (Uses)

(synergistic fungicidal composition)

IT 156052-68-5D, mixts. containing

RL: AGR (Agricultural use); BIOL (Biological study); USES (Uses)

(synergistic fungicidal compns.)

L3 ANSWER 19 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2000:335212 CAPLUS Full-text

DOCUMENT NUMBER: 132:339369

TITLE: An inhalation system containing a lipid

mixture  
 INVENTOR(S): Pilkiewicz, Frank G.  
 PATENT ASSIGNEE(S): USA  
 SOURCE: PCT Int. Appl., 36 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000027359	A1	20000518	WO 1999-US26858	
19991112 <--				
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
CA 2351063	A1	20000518	CA 1999-2351063	
19991112 <--				
EP 1128813	A1	20010905	EP 1999-958945	
19991112 <--				
EP 1128813	B1	20070214		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, CY				
HU 2001004255	A2	20020328	HU 2001-4255	
19991112 <--				
HU 2001004255	A3	20021228		
JP 2002529393	T	20020910	JP 2000-580590	
19991112 <--				
NZ 511568	A	20030829	NZ 1999-511568	
19991112 <--				
AU 766703	B2	20031023	AU 2000-16212	
19991112 <--				
AT 353630	T	20070315	AT 1999-958945	
19991112 <--				
ES 2281199	T3	20070916	ES 1999-958945	
19991112 <--				
EP 1839648	A2	20071003	EP 2007-1365	
19991112 <--				
EP 1839648	A3	20071121		
R: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
CN 100358494	C	20080102	CN 1999-815281	

19991112 <--  
     ZA 2001003645           A       20020805       ZA 2001-3645  
 20010504 <--  
     MX 2001004828           A       20020918       MX 2001-4828  
 20010511 <--  
     IN 2004DN03557          A       20050401       IN 2004-DN3557  
 20041211 <--  
 PRIORITY APPLN. INFO.:                           US 1998-108067P       P  
 19981112 <--                                       US 1998-108126P       P  
   EP 1999-958945       A3  
 19991112 <--                                       WO 1999-US26858       W  
 19991112 <--  
 AB    A system for administering a bioactive agent by inhalation  
       comprises a lipid mixture containing a phosphatidylcholine,  
       phosphatidylethanolamine, phosphatidylglycerol,  
       phosphatidylinositol, sterol, albumin and phosphatidic acid in  
       various combinations and ratios. The biol. active agent is a  
       drug, such as antitumor or antimicrobial agent, a compound  
       affecting endocrine function, an antibody, a gene, a cytokine, a  
       differentiating agent, etc.  
 TI    An inhalation system containing a lipid mixture  
 REFERENCE COUNT:           4       THERE ARE 4 CITED REFERENCES AVAILABLE  
 FOR THIS  
   RECORD. ALL CITATIONS AVAILABLE IN THE  
 RE FORMAT  
 PI    WO 2000027359 A1 20000518  
       PATENT NO.           KIND    DATE           APPLICATION NO.       DATE  
       -----  
 ----  
 PI    WO 2000027359           A1       20000518       WO 1999-US26858  
 19991112 <--  
     W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN,  
 CU, CZ,           DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,  
                   JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD,  
 MG, MK,           MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,  
 SL, TJ,           TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW  
     RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH,  
 CY, DE,           DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF,  
 BJ, CF,           CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG  
     CA 2351063           A1       20000518       CA 1999-2351063  
 19991112 <--  
     EP 1128813           A1       20010905       EP 1999-958945  
 19991112 <--  
     EP 1128813           B1       20070214  
     R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,  
 MC, PT,           IE, SI, LT, LV, FI, RO, CY  
     HU 2001004255           A2       20020328       HU 2001-4255

19991112 <--  
     HU 2001004255           A3     20021228  
     JP 2002529393           T     20020910     JP 2000-580590  
 19991112 <--  
     NZ 511568                A     20030829     NZ 1999-511568  
 19991112 <--  
     AU 766703                B2     20031023     AU 2000-16212  
 19991112 <--  
     AT 353630                T     20070315     AT 1999-958945  
 19991112 <--  
     ES 2281199               T3     20070916     ES 1999-958945  
 19991112 <--  
     EP 1839648               A2     20071003     EP 2007-1365  
 19991112 <--  
     EP 1839648               A3     20071121  
     R: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LI,  
 LU, MC,  
         NL, PT, SE  
     CN 100358494            C     20080102     CN 1999-815281  
 19991112 <--  
     ZA 2001003645           A     20020805     ZA 2001-3645  
 20010504 <--  
     MX 2001004828           A     20020918     MX 2001-4828  
 20010511 <--  
     IN 2004DN03557          A     20050401     IN 2004-DN3557  
 20041211 <--  
 PRAT US 1998-108067P       P     19981112 <--  
     US 1998-108126P       P     19981112 <--  
     EP 1999-958945       A3     19991112 <--


L3 ANSWER 20 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 1999:557667 CAPLUS Full-text  
 DOCUMENT NUMBER: 131:166517  
 TITLE: Synergistic fungicidal compositions  
           containing N-acetonylbenzamidates  
 INVENTOR(S): Young, David Hamilton; Wilson, Willie Joe;  
 Egan, Anne  
 PATENT ASSIGNEE(S): Ritchie; Michelotti, Enrique Luis  
 SOURCE: Rohm and Haas Company, USA  
           Eur. Pat. Appl., 15 pp.  
           CODEN: EPXXDW  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 4  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	----
EP 937396	A2	19990825	EP 1998-310539	
19981221 <--				
EP 937396	A3	19991208		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,				
MC, PT,				
IE, SI, LT, LV, FI, RO				



US 6004947	A	19991221	US 1998-148604
19980904 <--			
EP 1195089	A2	20020410	EP 2001-130309
19981221 <--			
EP 1195089	A3	20020424	
EP 1195089	B1	20031126	
R: DE, ES, FR, GB, IT			
EP 1247448	A2	20021009	EP 2002-15785
19981221 <--			
EP 1247448	A3	20021016	
EP 1247448	B1	20031210	
R: DE, ES, FR, GB, IT			
EP 1247449	A2	20021009	EP 2002-15786
19981221 <--			
EP 1247449	A3	20021016	
EP 1247449	B1	20040211	
R: DE, ES, FR, GB, IT			
EP 1247450	A2	20021009	EP 2002-15787
19981221 <--			
EP 1247450	A3	20021016	
EP 1247450	B1	20040107	
R: DE, ES, FR, GB, IT			
EP 1247451	A2	20021009	EP 2002-15788
19981221 <--			
EP 1247451	A3	20021016	
EP 1247451	B1	20031210	
R: DE, ES, FR, GB, IT			
ES 2207627	T3	20040601	ES 2002-15785
19981221 <--			
ES 2207628	T3	20040601	ES 2002-15788
19981221 <--			
ES 2210211	T3	20040701	ES 2002-15787
19981221 <--			
ES 2211845	T3	20040716	ES 2002-15786
19981221 <--			
AU 9912098	A	19990819	AU 1999-12098
19990114 <--			
AU 751144	B2	20020808	
TW 529907	B	20030501	TW 1999-88100522
19990114 <--			
CN 1229580	A	19990929	CN 1999-100316
19990120 <--			
CN 1128579	C	20031126	
MX 9900809	A	20000228	MX 1999-809
19990121 <--			
BR 9900173	A	20000502	BR 1999-173
19990126 <--			
JP 11310505	A	19991109	JP 1999-17816
19990127 <--			
US 6057356	A	20000502	US 1999-433974
19991104 <--			
US 6060490	A	20000509	US 1999-433973
19991104 <--			
MX 2002007916	A	20030425	MX 2002-7916
20020815 <--			
MX 2002007917	A	20030425	MX 2002-7917
20020815 <--			

MX 2002007918 A 20030425 MX 2002-7918  
 20020815 <--  
 MX 2002007921 A 20030425 MX 2002-7921  
 20020815 <--  
 MX 2002007922 A 20030425 MX 2002-7922  
 20020815 <--  
 PRIORITY APPLN. INFO.: US 1998-72725P P  
 19980127 <--  
 US 1998-148604 A  
 19980904 <--  
 EP 1998-310539 A3  
 19981221 <--  
 OTHER SOURCE(S): MARPAT 131:166517  
 AB Phytopathogenic fungi are controlled by the application of a selected fungicidally active N-acetonylbenzamide compound and a second fungicidally active compound selected from the group consisting of an inhibitor of respiration at cytochrome complex III, ziram, fluazinam, zarilamide, chlorothalonil, propamocarb, folpet, fosetyl-aluminum or a fungitoxic metabolite thereof, a triphenyltin-type fungicide and a copper-containing fungicide to plant seed, to plant foliage or to a plant growth medium. The compns. and method of use provide higher fungicidal activity than sep. use of the same compds. Cucumber downy mildew and tomato late blight were effectively controlled with different combinations of N-[3'-(1'-chloro-3'-methyl-2'-oxopentane)]-3,5-dichloro-4- methylbenzamide and propamocarb. Synergism was seen at the ratios: 1:42,1:21,1:10.5,1:5,1:2.5 and 1:1.  
 TI Synergistic fungicidal compositions containing N-acetonylbenzamides  
 L3 ANSWER 21 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 1984:97934 CAPLUS Full-text  
 DOCUMENT NUMBER: 100:97934  
 ORIGINAL REFERENCE NO.: 100:14785a  
 TITLE: Physiological and biochemical effects of  
 analogs of  
 the herbicide propyzamide  
 AUTHOR(S): Tissut, Michel; Aspe, Daniel; Meallier,  
 Pierre; Coste,  
 Camille  
 CORPORATE SOURCE: Lab. Physiol. Cell. Veg., Univ. Grenoble,  
 Saint-Martin-d'Herès, 38402, Fr.  
 SOURCE: Physiologie Vegetale (1983), 21(4), 689-99  
 CODEN: PHYVAP; ISSN: 0031-9368  
 DOCUMENT TYPE: Journal  
 LANGUAGE: French  
 GI



- AB The effects of the N-(1,1-dimethylpropynyl)benzamides (I, R = H, halo, CN, NO<sub>2</sub>, Me, and OMe) were studied on the photosynthesis of isolated chloroplasts, O consumption of isolated mitochondria, and growth of barley (*Hordeum vulgare*) seedlings. No correlation appeared between the effects on mitochondria or chloroplasts and I effect on barley seedlings. A 50% inhibition of photosynthesis was measured at the photosystem II level for concns. between 60  $\mu$ M and 0.3 mM. Inhibition of the mitochondrial electron flow appeared in the flavoprotein region for concns. 100  $\mu$ M and saturation Except for I (R = NO<sub>2</sub>), I produced the same herbicidal symptoms on barley seedlings. The concns. needed for such effects were between 1  $\mu$ M and 1 mM. The Cl at the 3 and 5 positions of the benzene ring greatly enhanced the mitosis inhibition in barley seedlings.
- TI Physiological and biochemical effects of analogs of the herbicide propyzamide
- TI Physiological and biochemical effects of analogs of the herbicide propyzamide
- SO Physiologie Vegetale (1983), 21(4), 689-99  
CODEN: PHYVAP; ISSN: 0031-9368
- AB The effects of the N-(1,1-dimethylpropynyl)benzamides (I, R = H, halo, CN, NO<sub>2</sub>, Me, and OMe) were studied on the photosynthesis of isolated chloroplasts, O consumption of isolated mitochondria, and growth of barley (*Hordeum vulgare*) seedlings. No correlation appeared between the effects on mitochondria or chloroplasts and I effect on barley seedlings. A 50% inhibition of photosynthesis was measured at the photosystem II level for concns. between 60  $\mu$ M and 0.3 mM. Inhibition of the mitochondrial electron flow appeared in the flavoprotein region for concns. 100  $\mu$ M and saturation Except for I (R = NO<sub>2</sub>), I produced the same herbicidal symptoms on barley seedlings. The concns. needed for such effects were between 1  $\mu$ M and 1 mM. The Cl at the 3 and 5 positions of the benzene ring greatly enhanced the mitosis inhibition in barley seedlings.
- ST barley photosynthesis respiration chlorodimethylpropynyl benzamide; propyzamide analog *Hordeum* growth
- IT Plant respiration  
(by barley mitochondria, dichloro(dimethylpropynyl)benzamide and its derivs. effect on)
- IT Photosynthesis  
Plant growth and development  
(by barley, dichloro(dimethylpropynyl)benzamide and its derivs. effect on)
- IT Barley  
(growth and photosynthesis and respiration by, dichloro(dimethylpropynyl)benzamide and its derivs. effect on)
- IT Chloroplast  
(photosynthesis of, by barley, dichloro(dimethylpropynyl)benzamide and its derivs. effect on)
- IT Mitochondria  
(respiration by, of barley, dichloro(dimethylpropynyl)benzamide and its derivs. effect on)
- IT 23950-58-5 23950-58-5D, derivs. 23955-55-7 24911-27-1

89026-33-5

89026-34-6 89026-35-7 89026-36-8 89026-37-9

RL: BIOL (Biological study)

(growth and photosynthesis and respiration by barley response to)

L3 ANSWER 22 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1972:33143 CAPLUS Full-text

DOCUMENT NUMBER: 76:33143

ORIGINAL REFERENCE NO.: 76:5389a,5392a

TITLE: Soil respiration and enzyme activities of herbicide-treated vineyard soils. III

AUTHOR(S): Walter, B.; Bastgen, D.

CORPORATE SOURCE: Abt. Bodenkd., Landes- Lehr- Versuchsanst. Trier,

Trier, Fed. Rep. Ger.

SOURCE: Weinberg & Keller (1971), 18(10), 465-74

CODEN: WBKRAC; ISSN: 0508-2404

DOCUMENT TYPE: Journal

LANGUAGE: German

AB In 2-year field expts. on Devonian slate and shell-lime soils the influence of various herbicides in pre- and postemergence treatment on the biol. activity of vineyard soils was investigated. After preemergence herbicide application a repeated soil cultivation was made. The herbicides used were dichlorobenzonitrile+dichloro-thiobenzamide, dichlobenil, simazine+amitrole+MCPA, atrazine+mecoprop, and diquat+paraquat. By using the air-dried soil fraction<2 mm soil respiration as well as the dehydrogenase, phosphatase, urease, glucosidase, and invertase activities were tested. CO<sub>2</sub> production was reduced after herbicide treatment. There was no difference between the 2 soils used. Generally, there was an increase in the enzyme activities. Soil cultivation was of importance for the activities as could be demonstrated for glucosidase and invertase.

L3 ANSWER 23 OF 23 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1961:93942 CAPLUS Full-text

DOCUMENT NUMBER: 55:93942

ORIGINAL REFERENCE NO.: 55:17739h-1,17740a

TITLE: Polarographic studies on the concentration of oxygen

in broth and oxygen uptake rate of mycelium in submerged fermentation of *Penicillium*

*chrysogenum*

AUTHOR(S): Gondhalekar, R. S.; Phadke, R. S.

CORPORATE SOURCE: Hindustan Antibiotics Ltd., Pimpri

SOURCE: Journal of Scientific & Industrial Research (1960), 19C, 183-6

CODEN: JSIRAC; ISSN: 0022-4456

DOCUMENT TYPE: Journal

LANGUAGE: Unavailable

AB The O levels in broth and O uptake rates of the mycelium from fermentations of different strains of *P. chrysogenum* are measured polarographically. The O levels in the fermentations of strains producing pelleted mycelium are lower than the strains giving

filamentous mycelium. The polarographic residual currents of the broth filtrates are abnormally high in fermentations with low yields.

TI Polarographic studies on the concentration of oxygen in broth and oxygen

uptake rate of mycelium in submerged fermentation of *Penicillium chrysogenum*

SO Journal of Scientific & Industrial Research (1960), 19C, 183-6  
CODEN: JSIRAC; ISSN: 0022-4456

IT Fermentation

(by *Penicillium chrysogenum*, O concentration and respiration during)

IT Fungicides or Fungistats

(sulfamic acid derivs. as)

IT 1227-29-8, Benzamide, 4,4'-dithiobis- 2527-57-3,  
Benzamide, 2,2'-dithiobis- 16624-71-8, Benzenesulfonamide,  
4,4'-dithiobis- 104997-15-1, Benzenesulfonamide, 2,2'-dithiobis-  
104997-16-2, Benzenesulfonamide, 3,3'-dithiobis- 107920-19-4,  
Benzamide, 3,3'-dithiobis- 109692-80-0, Benzenesulfonamide,  
2,2'-dithiobis[N-ethyl- 109692-81-1, Benzenesulfonamide,  
3,3'-dithiobis[N-ethyl- 109692-82-2, Benzenesulfonamide,  
4,4'-dithiobis[N-ethyl- 113926-47-9, Benzenesulfonanilide,  
3,3'-dithiobis- 113926-48-0, Benzenesulfonanilide, 4,4'-  
dithiobis-  
113926-94-6, Benzenesulfonanilide, 2,2'-dithiobis- 114160-45-1,  
Benzenesulfonamide, 2,2'-dithiobis[N-butyl- 114160-46-2,  
Benzenesulfonamide, 3,3'-dithiobis[N-butyl- 114160-47-3,  
Benzenesulfonamide, 4,4'-dithiobis[N-butyl- 114160-48-4,  
Benzenesulfonamide, 2,2'-dithiobis[N,N-diethyl- 114160-49-5,  
Benzenesulfonamide, 3,3'-dithiobis[N,N-diethyl- 114160-50-8,  
Benzenesulfonamide, 4,4'-dithiobis[N,N-diethyl-  
(bactericidal and fungicidal action of)  
IT 5329-14-6, Sulfamic acid  
(derivs., as bactericides and fungicides)